arXiv:1211.5933v1 [cs.DS] 26 Nov 2012

Interval Deletion is Fixed-Parameter Tractable*

Yixin Cao and Dániel Marx
Computer and Automation Research Institute
Hungarian Academy of Sciences (MTA SZTAKI)
yixin@sztaki.hu, dmarx@cs.bme.hu

November 27, 2012

Abstract

We study the minimum *interval deletion* problem that asks for the removal of a set of at most k vertices to make a graph of n vertices into an interval graph. We present a parameterized algorithm of runtime $10^k \cdot n^{\mathcal{O}(1)}$ for this problem, i.e., we show that the problem is fixed-parameter tractable.

1 Introduction

A graph is an interval graph if its vertices can be assigned to the intervals of the real line such that there is an edge between two vertices if and only if their corresponding intervals intersect. Interval graphs are the natural models in many applications, among which the most cited ones include DNA chains in biology [3], which actually initiated and motivated the study of interval graphs [17], and jobs scheduling in industrial engineering [2]. As induced cycles of more than 3 vertices (called *holes*) cannot be represented by intervals, an interval graph contains no hole. In other words, interval graphs are a subset of chordal graphs, which are exactly the set of hole-free graphs. This containment is proper as there are chordal graphs that cannot be represented by intervals. It is known that a chordal graph is an interval graph if and only if it contains no *asteroidal triple*, i.e. three vertices each pair of which is connected by a path avoiding neighbors of the third one [17].

The properties and the recognition of interval and chordal graphs are well understood; in particular, many NP-hard problems (coloring, maximum independent set, etc.) are known to be polynomial solvable when restricted to these classes of graphs. Therefore, one would like to generalize these results to graphs that do not belong to these classes, but close to them in the sense that have only a few "erroneous"/"missed" edges or vertices. For such generalizations, one would like to know how far the given graph is from the class and to find the erroneous/missed elements. This leads us naturally to the area of graph modification problems, where given a graph G, the task is to apply a minimum number of operations on G to make it a member of the class $\mathcal{F}$. Depending on the operations we allow, we can consider completion (edge addition), edge deletion, and vertex deletion versions of these problems. Many classical graph-theoretic problems can be formulated as graph deletion to special graph classes. For instance, VERTEX COVER, FEEDBACK VERTEX SET, CLUSTER VERTEX DELETION, and ODD CYCLE TRANSVERSAL problems can be viewed as a vertex deletion problem where the class $\mathcal{F}$ is the class of all empty graphs, trees, clusters, and bipartite graphs, respectively.

Unfortunately, most of these problems are NP-hard: for example, a classical result of Lewis and Yannakakis [18] shows that the vertex deletion problem is NP-hard for *every* nontrivial and hereditary class $\mathcal{F}$. Therefore, early work of Kaplan et al. [14] and Cai [6] focused on the fixed-parameter tractability of modification problems. Recall that a problem, parameterized by k, is *fixed-parameter tractable (FPT)* if there is an algorithm with runtime $f(k) \cdot n^{O(1)}$, where f is a computable function depending only on k [9]. Kaplan et al. [14] and Cai [6] observed that if a hereditary graph class $\mathcal{F}$ can be characterized by a finite number of forbidden (induced) subgraphs, then the fixed-parameter tractability of the modification problem follows from a basic bounded search tree algorithm. However, many important graph classes, such as trees, bipartite graphs, or chordal graphs have minimal obstructions of arbitrary large size (cycles, odd cycles, and holes, repsectively). It is much more challenging to obtain fixed-parameter tractability results for such

---

[*] Research supported by the European Research Council (ERC) grant "PARAMTIGHT: Parameterized complexity and the search for tight complexity results", reference 280152.

classes, see results, for example, on bipartite graphs [21, 15], planar graphs [20, 15], acyclic graphs [7], and minor-closed classes [1, 10].

Even though chordal graphs have minimal obstructions of unbounded size, CHORDAL COMPLETION (make the graph chordal by the addition of at most k edges) can still be solved by a bounded search tree algorithm by observing that a large hole immediately implies a negative answer to the problem [14, 6]. No such simple argument works for CHORDAL DELETION (make the graph chordal by removing at most k edges/vertices) and its fixed-parameter tractability was obtained by a completely different and much more complicated approach [19]. Observing the fixed-parameter tractability of CHORDAL COMPLETION, Kaplan et al. [14] asked if the apparently harder INTERVAL COMPLETION problem is FPT as well. Villanger et al. [22] resolved this question by giving a $k^{2k} \cdot n^{O(1)}$ time algorithm for INTERVAL COMPLETION. In this paper, we answer the complementary question about the vertex deletion version of the problem, resolving the perhaps most natural remaining open question in the area:

Theorem 1.1 (Main result). *There is a* $10^k \cdot n^{O(1)}$ *algorithm for deciding if there is a set of at most* k *vertices whose deletion makes the graph* G *an interval graph.*

Let us point out that, in a sense, INTERVAL DELETION is inherently harder than INTERVAL COMPLETION. The reason is that in the INTERVAL COMPLETION problem one can first use a simple bounded search tree algorithm to try every minimal way of completing all the holes; therefore, one can assume that the graph is chordal. On the other hand, the number of minimal ways of breaking the holes by removing vertices is not bounded by a constant and it is not clear how the very complex CHORDAL DELETION algorithm of [19] is of any help for INTERVAL DELETION. Indeed, we spend significant effort in the present paper to ensure that the graph is chordal. We overcome this difficulty by developing a reduction rule based on the modular decomposition of the graph and by analyzing the structural properties of reduced graphs. It turns out that the holes remaining in a reduced graph interact in a very special way (each hole is fully contained in the closed neighborhood of any other hole). This property allows us to prove that the number of minimal ways of breaking the holes is polynomially bounded, thus a simple branching step can reduce the problem to the case when the graph is chordal. As another consequence of our reduction rule, we can prove that this chordal graph already has a structure close to interval graphs (it has a clique tree that is a caterpillar). We can show that in such a chordal graph, asteriodal triples interact in a well-behaved way and we can find a set of 10 vertices such that there always exists a minimum solution that contains at least one of these 10 vertices. Therefore, we can complete our algorithm by branching on the deletion of one of these vertices.

2 Outline

The purpose of this section is to describe the main steps of our algorithm at a high level. We say that a set $Q \subset V(G)$ is an *interval deletion set* if $G - Q$ is an interval graph. An interval deletion set Q is *minimum* if there is no strictly smaller interval deletion set, and it is *minimal* if no proper subset $Q' \subset Q$ is an interval deletion set. A *minimal forbidden induced subgraph* is an induced subgraph $G[U]$ that is not an interval graph, but $G[U']$ is an interval graph for every proper subset $U' \subset U$. A *minimal forbidden set* is a set of vertices inducing a minimal forbidden induced subgraph. Clearly, set Q is an interval deletion set if and only if it intersects every minimal forbidden set. Our goal is to find an interval deletion set of size at most k. For technical reasons, it will be convenient to define the problem as follows:

INTERVAL DELETION: Given a graph G and an integer k, return

- if an interval deletion set of size $\leqslant k$ exists, a *minimum* interval deletion set $Q \subset V(G)$;
- if no interval deletion set of size $\leqslant k$ exists, "NO".

PHASE 1: Reductions. The first phase of the algorithm applies two reduction rules exhaustively. Each reduction rule either simplifies the instance or branches into a constant number of instances with strictly smaller parameter value. The first reduction rule is fairly simple: we destroy every forbidden induced subgraph of size at most 10.

Reduction 1. [Small Forbidden Induced Subgraphs] *Given an instance* (G, k) *and a minimal forbidden set* X *of no more than* 10 *vertices, we branch into* $|X|$ *instances,* $(G - v, k - 1)$ *for each* $v \in X$.

A graph on which Reduction 1 cannot be applied is called *prereduced*. By definition, each minimal forbidden induced subgraph that witnesses an asteroidal triple or hole in a prereduced graph contains at least 11 vertices. It can be checked in polynomial time whether Reduction 1 is applicable by enumerating every set X of size at most 10.

The second reduction rule is less obvious and more involved. Recall that a subset M of vertices forms a *module* if each vertex in M has the same neighbors out of M. A module M of G is *nontrivial* if $1 < |M| < |V(G)|$. We observe (see Section 4.2) that a minimal forbidden set X of at least 5 vertices is either fully contained in a module M or contains at most one vertex of M. Moreover, if $X \cap M = \{x\}$, then replacing x by any other vertex $x' \in M \setminus \{x\}$ in X results in another minimal forbidden set. This permits us to branch on modules, as described in the following rule.

Reduction 2. [Main] *Given an instance $I = (G, k)$ where the graph G is prereduced, and a nontrivial module M that does not induce a clique.*

1. *If every minimal forbidden set U is contained in M, then return the instance $(G[M], k)$.*
2. *If no minimal forbidden set is contained in M, then return the instance (G_M, k), where G_M is obtained from G by inserting edges to make $G[M]$ a clique.*
3. *Otherwise, we solve three instances: $I_1 = (G - M, k - |M|)$, $I_2 = (G[M], k - 1)$, and $I_3 = (G', k - 1)$, where G' is obtained from G by replacing M with a clique M' of $(k + 1)$ vertices; let Q_1, Q_2, Q_3 be the solutions of these instances, respectively. We return either $Q_1 \cup M$ or $Q_2 \cup Q_3$ ("NO" when $|Q_2 \cup Q_3| > k$), whichever is smaller.*

That is, in the third case we branch into two directions: the solution is obtained either as the union of M and the solution of I_1, or as the union of solutions of I_2 and I_3. The two branches correspond to the two cases where the solution fully contains M or only a minimum interval deletion set of $G[M]$ (i.e. Q_2), respectively. Note that in the the second branch, the vertices of the $(k + 1)$-clique M' are not part of Q_3, which is minimum and of size $\leqslant k$; hence $Q_2 \cup Q_3$ is indeed a subset of $V(G)$. Moreover, we have to clarify what the behavior of the reduction is if one of Q_1, Q_2, Q_3 is "NO." If Q_2 or Q_3 is "NO," then we define $Q_2 \cup Q_3$ to be "NO" as well. If one of Q_1 and $Q_2 \cup Q_3$ is "NO," we return the other one; if both of them are "NO," we return "NO" as well.

A graph on which Reductions 1 and 2 are not applicable is called *reduced*; in such a graph, every nontrivial module induces a clique. In Section 4, we prove the correctness of the reductions and that it can be checked in polynomial time if a reduction rule is applicable. Hence after exhaustive application of the reduction rules, we may assume that the problem instance is reduced.

The reductions are followed by a comprehensive study on reduced graphs that prove two crucial combinatorial statements. A vertex is called *simplicial* if its neighborhood induces a clique. The first combinatorial statement shows that every asteroidal triple has a simplicial vertex. In fact, we prove a slightly stronger statement, which shows that the so-called "shallow terminal" of the asteroidal triple is simplicial. Of an asteroidal triple $\{x, y, z\}$ witnessed by a minimal forbidden induced subgraph that is not a hole, we say that x is a shallow terminal if the defining path between y and z has length strictly larger than 3. As we shall see in Section 4.1, every minimal forbidden induced subgraph of size at least 11 and different from a hole has a shallow terminal.

Theorem 2.1. [Simplicial terminals] *Let G be a reduced graph, then all shallow terminals in G are simplicial.*

The way we show that a shallow terminal is simplicial is by showing that it is contained in a module whose neighborhood is a clique. If Reductions 2 cannot be applied, then this module induces a clique and every vertex in it is simplicial.

We say that two holes are *congenial* to each other if each hole is in the closed neighborhood of the other hole. It turns out that the holes are pairwise congenial in a reduced graph.

Theorem 2.2. [Congenial holes] *Let G be a reduced graph that is not chordal, then all holes of G are congenial to each other.*

PHASE 2: Breaking holes. Theorem 2.2 is of fundamental importance for the second phase, where we want to get rid of holes. Observe that Theorem 2.2 implies that if v is a vertex of a hole, then $N[v]$ intersects every hole and thus makes a hole cover. Intuitively, this suggests that a minimal hole cover has to be very local in a certain sense. Indeed, by relating minimal hole covers in the reduced graph and minimal separators in the subgraph $G - N[v]$, we are able to establish a quadratic bound on the number of minimal hole covers, and more importantly, a cubic time algorithm to construct them.

Theorem 2.3. [Hole covers] *Let G be a reduced graph. Then G contains at most n^2 minimal hole covers, and they can be found in $\mathcal{O}(n^3)$ time.*

Any interval deletion set must be a hole cover, and thus contain a minimal hole cover. This allows us to branch into at most n^2 instances, each of which has a chordal graph. Note that this branching is applied only once, hence only a polynomial factor will be induced.

PHASE 3: Asteroidal triples in a caterpillar decomposition. As all the holes have been broken, the graph is already chordal at the onset of the third phase, and it remains to destroy remaining asteroidal triples. Note that after removing some vertices, the graph remains prereduced and it is still true that every shallow terminal is simplicial (Theorem 2.1).[1] The removal of all simplicial vertices breaks all asteroidal triples, hence makes an interval graph. This implies that the chordal graph at this stage has a very special structure: it has a clique tree decomposition where the tree is a caterpillar, i.e., a path with degree-1 vertices attached to it. In other words, all those vertices that do not define any asteroidal triples can be arranged in a linear way, which greatly simplifies the examination of interactions between asteroidal triples. As a consequence, we can select an asteroidal triple that is minimal in a certain sense, and single out 10 vertices such that there must exist a minimum interval deletion set destroying this asteroidal triple with one of these 10 vertices. We can now safely branch on removing one of these 10 vertices.

Theorem 2.4. [ATs in chordal graphs] *There is a $10^k \cdot n^{O(1)}$ time algorithm for the* INTERVAL DELETION *problem when the input graph is prereduced, chordal, and every shallow terminal is simplicial.*

Putting together these steps, the fixed-parameter tractability of INTERVAL DELETION follows (see Figure 1).

```
Algorithm Interval-Deletion($G, k$)
INPUT: a non-interval graph $G$ and a positive integer $k$
OUTPUT: a minimum interval deletion set $Q \subset V(G)$ of size $\leqslant k$ or "NO".
1    Reduction 1: Let $U$ be a minimal forbidden induced subgraph of at most 10 vertices;
         branch on deleting one vertex of $U$.
\\ the graph will then be prereduced and remains so henceforth;
2    Reduction 2: Let $M$ be a nontrivial module of $G$ not inducing a clique.
2.1      if all minimal forbidden sets of $G$ are contained in $M$ then
             return Interval-Deletion($G[M], k$).
2.2      else if no minimal forbidden set is contained in $M$ then
             return Interval-Deletion($G', k$), where edges are inserted to make $G[M]$ a clique.
2.3      else branch into three instances $I_1$, $I_2$, $I_3$;
\\ the graph will then be reduced;
3    Use the algorithm of Theorem 2.3 to enumerate the at most $n^2$ minimal hole covers of $G$.
\\ the graph might not remain reduced, but is prereduced and chordal, and every shallow terminal is simplicial;
4    for each minimal hole cover $HC$ do
4.1      Use the algorithm of Theorem 2.4 to solve $(G - HC, k - |HC|)$;
5    return the smallest solution obtained, or "NO" if all solutions are "NO".
```

Figure 1: Outline of algorithm for INTERVAL DELETION

Proof (of Theorem 1.1). The algorithm described in Figure 1 solves the problem by making recursive calls to itself, or calling the algorithm of Theorem 2.4 $\mathcal{O}(n^2)$ times. In the former case, at most 10 recursive calls are made, all with parameter value at most $k - 1$. In the latter case, the running time is $10^k \cdot n^{O(1)}$. It follows that the total running time of the algorithm is $10^k \cdot n^{O(1)}$. □

The paper is organized as follows. Section 3 sets the definitions and reviews some basic facts. Section 4 presents the details of the first phase. The next four sections are devoted to demonstrate the correctness of Theorems 2.1–2.4. Sections 5 and 6 put shallow terminals and congenial holes under thorough examination, and prove combinatorial statements Theorems 2.1 and 2.2, respectively. Section 7 proves Theorem 2.3 (the bound on the minimal hole covers) and Section 8 presents the algorithm that destroys asteroidal triples and proves Theorem 2.4. Section 9 closes this paper by some possible improvement and new directions.

DISCLAIMER. Independent of our work, Arash Rafiey and some anonymous author(s) claim a similar result with an approach different from ours (available at: `http://folk.uib.no/ara099/publication/edge_vertex.pdf`).

---

[1] However, the graph might no longer be reduced, as new nontrivial non-clique modules can appear. We could run the reduction of Phase 1 again to obtain a reduced instance, but there is no reason to do so at this point.

We both start from graphs without small forbidden induced subgraphs, however our main steps divert thereafter. We start with a rather simple reduction and perform a deep combinatorial study of reduced graphs. This allows us to get rid of holes first, and then we focus on the remaining asteroidal triples. Their work handles asteroidal triple first, and then holes. In particular, it claims to be able to locate a special asteroidal triple in a greedy way to branch on and then to break holes that are found also in a greedy way, but we have encountered some difficulties in fully verifying the correctness of these steps.

3 Preliminaries

All graphs discussed in this paper shall always be undirected and simple. A graph is given by its vertex set V and edge set E, whose sizes are $n = |V|$ and $m = |E|$ respectively. If a pair of vertices v_1 and v_2 are connected by an edge, they are *adjacent* to each other, and denoted by $v_1 \sim v_2$, otherwise *non-adjacent* and denoted by $v_1 \not\sim v_2$. By $v \sim X$ we mean v is adjacent to at least one vertex of the set X. Two vertex sets X and Y are completely connected if $x \sim y$ for each pair of $x \in X$ and $y \in Y$. A graph is *complete* if each pair of vertices are adjacent. A *clique* in a graph is a subgraph that is complete, and a clique is maximal if it is not contained in another clique. A neighbor of a vertex is another vertex that is adjacent to it, and the set of *neighborhood* of a vertex v is denote by $N(v)$. The *closed neighborhood* of v is defined as $N[v] = N(v) \cup \{v\}$. This is generalized to a vertex set U, whose closed neighborhood and neighborhood are defined to be $N[U] = \bigcup_{v \in U} N[v]$ and $N(U) = N[U] \backslash U$. The $N_U(v)$ ($N_U[v]$) stands for the neighbors of v in the set U, i.e. $N_U(v) = N(v) \cap U$ ($N_U[v] = N(v) \cap U$), and it is immaterial whether $v \in U$ or not. A vertex is *simplicial* if its neighbors induce a clique. The subgraph of a graph G induced by a subset of vertices U is denoted by $G[U]$, and $G - U$ is used as a shorthand for $G[V \setminus U]$. All subgraphs mentioned in this paper are induced; hence a subgraph G' of G is *proper* if and only if $V(G') \subset V(G)$.

A sequence of *distinct* vertices $v_0 v_1 \dots v_l$ such that $v_i \sim v_{i+1}$ for each $0 \leqslant i < l$ is called a *path*, or a v_0-v_l *path* when the two end vertices are of special interest, and its *length* is defined to be l. If $l > 1$ and $v_0 \sim v_l$, then the sequence $v_0 v_1 \dots v_l v_0$ is called a *cycle*, whose *length* is defined to be $l + 1$. As an abuse of notation, by $u \in P$ (or $u \in C$) we mean that the vertex u appears in the path P (or cycle C), i.e. we also consider a path (or cycle) as the set of elements in the sequence. A *chord* in a path or cycle is an edge between two non-consecutive vertices in the path or cycle (note that the edge $v_0 v_l$, if exists, is a chord in the path $v_0 v_1 \dots v_l$, but not in the cycle $v_0 v_1 \dots v_l v_0$). A chordless cycle of length at least 4 is called a *hole*. It is easy to verify that no shortest path can contain a chord, so between each pair of vertices of a connected graph there is a chordless path. A graph is *chordal* if any cycle of length at least 4 contains a chord, in other words, the graph contains no hole.

Chordal graphs enjoy several important and related characterizations. A set of vertices S is an x-y *separator* if there is no x-y path in the subgraph $G - S$, and *minimal* x-y *separator* if no proper subset of S separates x and y. For any pair of vertices x and y, a minimal x-y separator is also called a *minimal separator*. A graph is chordal if and only if each minimal separator in it is a clique [8]. A nontrivial chordal graph contains at least two simplicial vertices, and there is at least one simplicial vertex in each connected component after the removal of any separator.

A tree $\mathcal{T}$ whose nodes are the maximal cliques of a graph G is a *(maximal) clique tree* of G if it satisfies the following conditions: any two adjacent nodes K_i and K_j defines a minimal separator that is $K_i \cap K_j$; for any vertex $x \in V$, the cliques containing x correspond to a subtree of $\mathcal{T}$. A graph is chordal if and only if it has such a clique tree. A clique tree of a graph G will be denoted by $\mathcal{T}(G)$, and when the graph G is clear from the context, we use $\mathcal{T}$. Without distinguishing the node in a clique tree and the maximal clique in the graph G corresponding to it, we use K to denote both. A set of vertices is a minimal separator of G if and only if it is the common vertices of $K_i \cap K_j$, denoted by $S_{i,j}$, for an edge $K_i K_j$ in $\mathcal{T}$ [5]. Moreover, $K_i \cap K_j$ for an edge $K_i K_j$ in $\mathcal{T}$ is a minimal x-y separator for any $x \in K_i \setminus K_j$ and $y \in K_j \setminus K_i$. Since there are at most n maximal cliques in a chordal graph of n vertices [8], a clique tree $\mathcal{T}$ is simpler than G, and commonly called a compact representation of G.

As an interval graph is chordal, all the above properties also apply to interval graphs. Moreover, the clique tree of interval graphs admit one more characterization, which implies that the clique tree of an interval graph is always a path..

Theorem 3.1 ([11]). *A graph G is an interval graph if and only if the maximal cliques of G can be linearly ordered such that, for each vertex v, the maximal cliques containing v occur consecutively.*

We direct the reader to [4, 12] for more detailed studies on chordal graphs and interval graphs.

4 Reduction rules and branching

This section discusses the reduction rules described in Section 2 in more detail. Section 4.1 provides more details about Reduction 1 (removing small forbidden induced subgraphs), while Section 4.2 proves the correctness of Reduction 2 (reduction on modules).

4.1 Forbidden induced subgraphs

Three vertices form an *asteroidal triple* if each pair of them is connected by a path that avoids the neighbors of the third vertex. As an abuse of notation, we also use asteroidal triple (AT) to refer to the subgraph induced by union of these three paths, and the three defining vertices will be called *terminals* of this AT. By definition, the terminals are non-adjacent. Lekkerkerker and Boland observed that a graph is an interval graph if and only if it is chordal and contains no AT [17]. Not stopping here, they rolled up their sleeves and got their hands dirty by checking each possible forbidden induced subgraph. Their effort brought the following less beautiful but more useful characterization, here a minimal non-interval graph refers to a graph G whose every proper induced subgraph is an interval graph but itself is not.

Theorem 4.1 ([17]). *A minimal non-interval graph is either a hole or an asteroidal triple depicted in Figure 2.*



(a) long claw (b) whipping top (c) net (d) tent



(e) †-AT $(s : c : l, B, r)$ $(|B| \geqslant 3)$ (f) ‡-AT $(s : c_1, c_2 : l, B, r)$ $(|B| \geqslant 2)$

Figure 2: Minimal asteroidal triples in a chordal graph (terminals are marked as squares).

Some remarks are in order. First, it is easy to verify (by picking any three non-adjacent vertices from it as terminals) that a hole of 6 or more vertices satisfies the definition of AT, but following convention, we only refer it as a hole, and reserve the term AT for graphs listed in Figure 2. Second, the set of ATs depicted in Figure 2 are not a literal copy of Lekkerkerker and Boland's original list [17]. More specifically, a net is actually a †-AT with $d = 2$, and a tent is actually a ‡-AT with $d = 1$. We single out the net and tent for the convenience of later presentation. To avoid ambiguities, in this paper we explicitly require the length of the longest defining path of a †-AT and a ‡-AT to be at least 4 ($d \geqslant 3$) and 3 ($d \geqslant 2$) respectively. Third, each of the four subgraphs at the first row consists of constant number (6 or 7) of vertices, and thus can be easily located and disposed of by standard enumeration[2]. For the purpose of the current paper, we are mainly concerned with the two kinds of ATs at the second row, whose sizes are unbounded. In the three paths defining a †-AT (‡-AT resp.), two of them have length exactly 3 (2 resp.), and the other strictly larger than 3 (2 resp.). Among the three terminals, the one of distance 3 (a †-AT) or 2 (a ‡-AT) to both other terminals is called the *shallow terminal*, whose neighbor(s) are the *center(s)*. The other two terminals, i.e. ends of the long defining path (length > 3), are called *base terminals*. All other vertices form the *base*, which is a sub-path of the long defining path. We use $(s : c : l, B, r)$ ($(s : c_1, c_2 : l, B, r)$ resp.) to denote the †-AT (‡-AT resp.) with shallow terminal s, center c (centers c_1 and c_2 resp.), base terminals l, r, and base $B = \{b_1, \dots, b_d\}$. As a notational convenience, we will also use b_0 and b_{d+1} to refer to the base terminals l and r, respectively. The center(s) and vertices in the base are called non-terminal vertices.

Observe that none nonterminal vertex of the graphs appearing in Figure 2 is simplicial, and holes do not have simplicial vertices either:

---

[2] The whipping top is also kind of special, in the sense that its three terminals are not symmetric. Also, with one edge removed it contains a net as a proper subgraph, and with two edges it turns into a †-AT with $d = 3$.

Proposition 4.2. *No non-terminal vertex of an AT can be simplicial.*

A small forbidden induced subgraph is a small asteroidal triple T or a short hole H that consists of at most 10 vertices. In $\mathcal{O}(n^4)$ time we can find a small forbidden induced subgraph subgraph by checking each three vertices to see they are three consecutive vertices of a hole, or the terminals of an AT, and finding the path(s) accordingly. A graph on which Reduction 1 is not applicable is called *prereduced*. Hence any AT or hole in a prereduced graph contains at least 11 vertices, and in particular it cannot be a long claw, whipping top, net, or tent. The base of a †-AT (‡-AT resp.) in a prereduced graph contains at least 9 (8 resp.) vertices.

The purpose of the following two propositions and a detailed proof is twofold. This special structure arises frequently in this paper and we do not want to repeat it again and again. This argument is exemplary in the sense that, by and large, most proofs of this paper exploit a similar contradictory arguments: They explicitly construct a forbidden induced subgraph, either a small AT or a short hole, assuming the property under discussion does not hold; because all graphs discussed henceforth are prereduced, such a contradiction will suffice to prove the desired property.

Proposition 4.3. *Let* $P = (v_0 \dots v_p)$ *be a chordless path of length* $p \geqslant 4$ *in a prereduced graph* G*, and* u *is a adjacent to every vertex of* P*. Then* $N[v_\ell] \subseteq N[u]$ *for each* $2 \leqslant \ell \leqslant p-2$.

Proof. Assume to the contrary, there is a vertex $x \in N[v_\ell] \setminus N[u]$, then we show the existence of a short hole or small AT in G, thus contradicting the assumption that G is prereduced. Note that $x \not\sim v_i$ for any $i \leqslant \ell-2$ or $i \geqslant \ell+2$, as otherwise, there is a 4-hole $(uv_ixv_\ell u)$ (here $v_i \not\sim v_\ell$ because P is chordless). There is • a 4-hole $(uv_{\ell-1}xv_{\ell+1}u)$ if x is also adjacent to both $v_{\ell-1}$ and $v_{\ell+1}$; • a tent $\{u, v_{\ell-1}, v_\ell, x, v_{\ell+1}, v_{\ell+2}\}$ if x is adjacent to $v_{\ell+1}$ but not $v_{\ell-1}$; • a tent $\{u, v_{\ell-2}, v_{\ell-1}, x, v_\ell, v_{\ell+1}\}$ if x is adjacent to $v_{\ell-1}$ but not $v_{\ell+1}$; or • a whipping top $\{x, u, v_{\ell-2}, v_{\ell-1}, v_\ell, v_{\ell+1}, v_{\ell+2}\}$ otherwise (x is only adjacent to v_ℓ in the path). ☐

Proposition 4.4. *Let* $P = (v_0 \dots v_p)$ *be a chordless path of length* $p \geqslant 3$ *in a prereduced graph* G*, and* u *is a adjacent to all vertices in* P*. Then* $N[v_\ell] \cap N[v_{\ell+1}] \subseteq N[u]$ *for each* $1 \leqslant \ell \leqslant p-2$.

Proof. There is a tent $\{u, v_{\ell-1}, v_\ell, x, v_{\ell+1}, v_{\ell+2}\}$ if $x \in N[v_\ell] \cap N[v_{\ell+1}] \setminus N[u]$. ☐

Given a non-empty set X of vertices, we denote by $\widehat{N}(X)$ the set of *common neighbors* of X, that is, $v \in \widehat{N}(X)$ if and only if $v \sim x$ for each vertex $x \in X$. It is easy to verify that in a prereduced graph, if X itself does not induce a clique, then $\widehat{N}(X)$ does, as otherwise two non-adjacent vertices in $\widehat{N}(X)$, together with two non-adjacent vertices in X, will induce a 4-hole. In particular, we have the following proposition. We remark that the simple fact that common neighbors of two non-adjacent vertices in prereduced graphs must be adjacent will be frequently used is many proofs.

Proposition 4.5. *Let* X *be a set of vertices of a prereduced graph that induces either a hole, an AT, or a path of length at least* 2*. Then* $\widehat{N}(X)$ *induces a clique.*

4.2 Modular decomposition

A subset of vertices $M \subseteq V(G)$ form a *module* of G if each vertex in M has the same neighbors out of M. In other words, for any pair of vertices $u, v \in M$ and vertex $x \notin M$, $u \sim x$ if and only if $v \sim x$. The set $V(G)$ and each set of a single vertex are always modules; we call such modules *trivial*. A brief inspection of Figure 2 shows that a minimal AT does not have any nontrivial modules and holes of length 4 are the only holes having nontrivial modules. Thus we can derive:

Proposition 4.6. *Let* M *be a module, and* U *be a minimal forbidden set with* $|U| > 4$*. Then either* $U \subseteq M$*, or* $|M \cap U| \leqslant 1$.

This observation allows us to prove the following statement, which is the main combinatorial reason behind the correctness of the branching in Reduction 2:

Theorem 4.7. *Let* G *by a graph that contains no* 4*-hole and* M *be a module in* G*. A minimum interval deletion set of* G *contains either all vertices of* M*, or only a minimum interval deletion set to* $G[M]$.

Proof. Let Q be a minimum interval deletion set of G such that $M \not\subseteq Q$, otherwise we are already done. To show $Q_M = Q \cap M$ is precisely a minimum interval deletion set to G[M], it suffices to show that for any minimum interval deletion set Q'_M of G[M], the set $Q' = (Q \setminus Q_M) \cup Q'_M$ is an interval deletion set of G: Trivially Q_M is an interval deletion set of G[M]; if it is not minimum, then $|Q_M| > |Q'_M|$, and $|Q| > |Q'|$, which contradicts the fact that Q is minimum.

Consider a minimal forbidden set U in $G - Q'$; as there are no 4-holes by assumption, we have $|U| > 4$. By construction, the set Q'_M hits all minimal forbidden sets in $G[M]$, while $(Q \setminus Q_M)$ hits all minimal forbidden sets in $G - M$. Thus U contains a vertex $x \in M$, but it is not fully contained in M. By Lemma 4.6, x is the only vertex of U in M, and if $x' \in M \setminus Q$ (which exists by assumption), then $U' = U \setminus \{x\} \cup \{x'\}$ is also a minimal forbidden set: as $G[U]$ and $G[U']$ are isomorphic. The set U' nonetheless is contained in $G - Q$, which contradicts that Q is a solution. Therefore, Q' must be a solution of G. ☐

For the exhaustive application of Reduction 2, we have to be able to find a nontrivial module that is not a clique.

Lemma 4.8. *We can find a nontrivial module M in polynomial time such that $G[M]$ is not a clique, if such a module exists.*

Proof. The algorithm described in Figure 3 finds such a module in a greedy manner. It starts from a pair of nonadjacent vertices u and v, and generates the module by adding vertices. Note that a vertex set is returned by this algorithm only when both $X = \emptyset$ and $M \neq V(G)$ are true, so M must be a nontrivial module. Now it remains to make sure that it always finds a nontrivial module if such a module U exists. By assumption that $G[U]$ is not a clique, there are a pair of nonadjacent vertices $u, v \in U$. We claim no vertex of $V(G) \setminus U$ will be added in this algorithm. Suppose the contrary and $x \in V(G) \setminus U$ be the first vertex added in some iteration. However, we should have $M \subseteq U$ at that beginning of this iteration, which implies $N_M(x)$ is either M or $\emptyset$, and thus $x \notin X$. Therefore, we end with a contradiction. ☐

```
for each pair of non-adjacent vertices $u$ and $v$ do
    $M = \{u, v\}$;
    while $M \neq V(G)$ do
        $X = \{x \notin M | 0 < |N_M(x)| < |M|\}$;
        if $X = \emptyset$ then return $M$;
        else $M = M \cup X$.
```

Figure 3: Algorithm that finds a nontrivial non-clique module

Indeed, one can easily verify that the module found by this algorithm is the minimum nontrivial one that contains both u and v. We are now ready to explain the application of Reduction 2 and prove its correctness.

Lemma 4.9. *Reduction 2 is correct and it can be checked in polynomial time whether Reduction 2 (and which case of it) is applicable.*

Proof. To see whether Reduction 2 is applicable or not, we use Lemma 4.8 to find a nontrivial module M that does not induce a clique. Now we check the conditions in order to figure out which case should apply. To check whether Case 1 holds, we need to check if there is a minimal forbidden set U not contained in M. By Proposition 4.6, such an U, if exists, contains at most one vertex x from M; and x can be replaced by any other vertex of M. Therefore, it suffices to pick any vertex $x \in M$, and test in linear time whether $G - (M \setminus x)$ is an interval graph. If it is not an interval graph, then there is an minimal forbidden set U not contained in M (as it contains at most one vertex of M). Otherwise, $G - (M \setminus x)$ is an interval graph for every $x \in M$, and there is no such U; hence Case 1 holds. To check whether Case 2 holds, observe that the condition "there is no minimal forbidden set contained in M" is equivalent to saying that $G[M]$ is an interval graph, which can be checked in linear time. In all remaining cases, we are in Case 3.

The correctness of the reduction is clear in Case 1: removing the vertices of $V(G) \setminus M$ does not make the problem any easier, as these vertices are not used by the minimal forbidden induced subgraphs.

In Case 2, the correctness of the reduction follows from the fact that G and G_M have the same set of minimal forbidden sets. Indeed, M induces an interval graph, thus if U is a minimal forbidden set in G or G_M, then Proposition 4.6 implies that U contains at most one vertex of M (observe that making M a clique does not introduce any new 4-holes, thus Proposition 4.6 is applicable for G_M). Therefore, the removal and addition of edges have no effect on the minimal forbidden sets, which means that Q is an interval deletion set of G if and only if it is an interval deletion set of G_M.

The correctness of Case 3 can be argued using Theorem 4.7. As the graph is prereduced, it does not contain any 4-hole, thus Theorem 4.7 implies that there is a solution containing all of M or its intersection with M is a minimum interval deletion set of $G[M]$. The two branches correspond to these two cases. The first branch is straightforward: we simply remove all vertices of M from the graph and solve the instance $I_1 = (G - M, k - |M|)$. It is the second branch (where we assume $M \nsubseteq Q$) that needs more explanation.

Recall that $I_2 = (G[M], k-1)$ and $I_3 = (G', k-1)$, where G' is obtained by removing M, adding a clique M' of size $k+1$, and connecting M' with every vertex of $N(M)$. Clearly M' is a module and induces an interval graph. We make the size of M' to be $k+1$ in order to "forbid" vertices of it to be picked as a part of solution Q_3.[3] It is clear that either solution Q_2 or Q_3 is "NO" will rule out the existence of an interval deletion set of G that does not fully contain M. In the rest of the proof, we show that the solution $Q_2 \cup Q_3$ returned by the algorithm, if not "NO", is indeed a minimum interval deletion set that does not fully include M.

We claim that if Q_2 and Q_3 are minimum interval deletion sets of I_2 and I_3, respectively, then $Q = Q_2 \cup Q_3$ is an interval deletion set of (G, k). Suppose that there is a minimal forbidden set U disjoint from Q. It cannot be fully contained in M, as $Q_2 \subseteq Q$ is an interval deletion set of $G[M]$. Then by Proposition 4.6, U contains exactly one vertex x of M and $U' = U \setminus \{x\} \cup \{x'\}$ is also a minimal forbidden set of G' for any $x' \in M'$. As Q_3 is an interval deletion set of G' disjoint from M', it has to contain a vertex of $U' \setminus \{x'\} = U \setminus \{x\}$. Now it remains to show $Q \subset V(G)$. Note that no minimal interval deletion set can intersect the $(k+1)$-clique in G' (according to Theorem 4.7, such a solution, if exists, must contain all vertices of the $(k+1)$-clique, and then has size strictly larger than k), thus Q_3 is indeed a subset of $V(G)$.

Having shown that the union of an interval deletion set of I_2 and an interval deletion set of I_3 is an interval deletion set of G, next we show that it is not smaller than the smallest interval deletion set Q' satisfying $M \not\subseteq Q'$. Suppose that Q' is an interval deletion set of G of size at most k with $M \not\subseteq Q'$; let $Q'_2 = Q' \cap M$ and $Q'_3 = Q' \setminus M$. We claim that Q'_2 and Q'_3 are interval deletion sets of I_2 and I_3, respectively. First, we argue that Q'_2 and Q'_3 are not empty; hence both of them has size at most $k-1$. The assumption that $G[M]$ is not an interval graph implies $Q'_2 \neq \emptyset$. By assumption, $M \not\subseteq Q'$, thus there is a vertex $x \in M \setminus Q'$. Now $Q'_3 = \emptyset$ would imply that $G - (M \setminus \{x\})$ is an interval graph, that is, there is no minimal forbidden set containing only one vertex of M, and it follows that we should have been in Case 1. Since $|Q'_2| \leqslant k-1$, it is clear that Q'_2 is a solution of instance $I_2 = (G[M], k-1)$. The only way Q'_3 is not a solution of I_3 is that there is a minimal forbidden set U containing a vertex of the $(k+1)$-clique introduced to replace M. As this $(k+1)$-clique is a module, Proposition 4.6 implies that U contains exactly one vertex y of this clique. But in this case $U' = U \setminus \{y\} \cup \{x\}$ (where x is a vertex of $M \setminus Q'$) is a minimal forbidden set disjoint from Q', a contradiction. Thus following from the fact that both Q_2 and Q_3 are minimum, the size of the interval deletion set Q returned by this branch is no more than Q'. ☐

5 Shallow terminals

This section proves Theorem 2.1. We would like to point out the resemblance between the statements here and Observations 5.2-5.5 Villanger et al. made on chordal graphs [22], by which our observation was inspired. Nonetheless, here we are not equipped with chordality, and thus their simple argument via a classic characterization of chordal graphs (each minimal separator is a clique) does not work. Our approach here is to show that each shallow terminal is contained in a nontrivial module whose neighborhood induces a clique; as Reduction 2 cannot be applied, all vertices in this module are simplicial.

Recall that an AT in a prereduced graph is either a †-AT or a ‡-AT. Let us start from a thorough scrutiny of neighbors of a shallow terminal.

Lemma 5.1. *Let* T *be a †-AT* $(s : c : l, B, r)$ *or ‡-AT* $(s : c_1, c_2 : l, B, r)$ *in a prereduced graph* G*, and* $x \in N(s)$.

(1) *Then* x *is also adjacent to the center(s) of* T *(different from* x*).*

(2) *Classifying* x *with respect to its adjacency to* B*, we have the following categories:*

(full) x *is adjacent to every vertex of* B.
Then x *is also adjacent to every vertex of* $N(s) \setminus \{x\}$.

(partial) x *is adjacent to some, but not all vertices of* B.
Then there is an AT T' *whose shallow terminal is* s*, one center is* x*, and base* B' *is a proper sub-path of* B.

(none) x *is adjacent to no vertex of* B.
Then $x \not\sim l$ *and* $x \not\sim r$ *hold as well and thus replacing the shallow terminal of* T *by* x *makes another AT.*

---

[3] Indeed, $\min(k+1, |N(M)|)$ vertices will suffice for our bookkeeping purpose, and an alternative way to is to add only one vertex but mark it as "forbidden".

Proof. Assume to the contrary, $x \not\sim c$ if T is a †-AT or (without loss of generality,) $x \not\sim c_2$ if T is a ‡-AT. We separate the proof of the first assertion based on $x \sim B$ or not.

$x \sim B$. (See Figure 6(a).) There is • a 4-hole $(xscb_ix)$ if T is a †-AT and $x \sim b_i$ for some $1 \leqslant i \leqslant d$; • a 4-hole (xsc_2b_ix) if T is a ‡-AT and $x \sim b_i$ for some $1 \leqslant i \leqslant d+1$;

$x \not\sim B$. (See Figure 6(b,c,d,e).) There is • a 5-hole $(xscb_1lx)$ or $(xscb_drx)$ if T is a †-AT, and $x \sim l$ or $x \sim r$, respectively; or • a 5-hole (xsc_2b_1lx) or 4-hole (xsc_2rx) if T is a ‡-AT, and $x \sim l$ or $x \sim r$, respectively. • a long-claw $\{x, s, c, b_1, l, b_d, r\}$ if T is a †-AT and $x \not\sim l, r$; • a net $\{x, s, l, c_1, r, c_2\}$ if T is a ‡-AT and $x \not\sim c_1, l, r$; or • a whipping top $\{r, c_2, s, x, c_1, l, b_1\}$ centered at c_2 if T is a ‡-AT and $x \not\sim l, r$, but $x \sim c_1$.

Neither of these cases is possible, and thus statement (1) is proved. For statement (2), let us handle the category "none" first. Note that x cannot be a center of T, as it is not adjacent to B. If $x \sim l$, then (xcb_1lx) is a 4-hole if T is a †-AT and (xc_2b_1lx) is a 4-hole if T is a ‡-AT. As a symmetrical argument shows that $x \not\sim r$. Now that x is adjacent to the center(s) of T and $x \not\sim B \cup \{l, r\}$, then by definition, $(x : c : l, B, r)$ ($(x : c_1, c_2 : l, B, r)$ resp.) makes another †-AT (‡-AT resp.)

Suppose now that x is in the category "full." Suppose the contrary and $x \not\sim v$ for some $v \in N(s) \setminus \{x\}$. We have already proved in statement (1) that v and x are adjacent to the center(s) of T (different from them). In particular, if one of v and x is a center, then they are adjacent. Therefore, we can assume that v and x are not centers. If $v \sim B$, that is, $x \sim b_i$ for some $1 \leqslant i \leqslant d$, then there is a 4-hole $(xsvb_ix)$. Otherwise, if $v \not\sim B$, then v is in the category "none" and we have already proved that such a vertex is not adjacent to l and r. When T is a †-AT, there is (see the first row of Figure 7) • a whipping top $\{v, c, l, b_1, x, b_d, r\}$ centered at c if $x \sim l, r$; • a net $\{v, c, l, b_1, r, x\}$ if $x \sim r$ but $x \not\sim l$ (similarly for $x \sim l$ but $x \not\sim r$); or • a †-AT $\{v : c : l, b_1, x, b_d, r\}$ if $x \not\sim l, r$. When T is a ‡-AT, there is (see the second row of Figure 7) • a tent $\{x, c_1, b_1, v, b_d, c_2\}$ if $x \sim l, r$; • a ‡-AT $(v : c_1, c_2 : l, b_1x, r)$ if $x \sim r$ but $x \not\sim l$ (similarly for $x \sim l$ but $x \not\sim r$); or • a ‡-AT $(v : c_1, c_2 : l, b_1xb_d, r)$ if $x \not\sim l, r$.

Finally, suppose that x is in category "partial," that is, $x \sim B$, but $x \not\sim b_i$ for some $1 \leqslant i \leqslant d$. In this case, we construct the claimed AT as follows. As the case $x \not\sim l$ but $x \sim r$ is symmetric to $x \sim l$ but $x \not\sim r$, it is ignored in the following, i.e. $x \sim r$ only if $x \sim l$. Let p be the smallest index such that $x \sim b_p$, and q be the smallest index such that $p < q \leqslant d+1$ and $x \not\sim b_q$ (q exists by assumptions). See Figure 8[4] and Table 1 for the structures for †-AT and ‡-AT respectively.

Table I. $T(s : c : l, r)$ is a †-AT.

	$q = p+1$	$q = p+2$	$q > p+2$
$p = 0$ (Fig.8a)	4-hole (xcb_1lx)	tent $\{l, x, s, c, b_2, b_1\}$	‡-AT $T(s : x, c : l, b_1 \ldots b_{q-1}, b_q)^2$
$p = 1$ (Fig.8b)	whipping top^1 $\{l, b_1, x, s, c, b_3, b_2\}$	net $\{l, b_1, s, x, b_3, b_2\}$	†-AT $T(s : x : l, b_1 \ldots b_{q-1}, b_q)^2$
$p > 1$ (Fig.8c)	long-claw^1 $\{b_{p-2}, b_{p-1}, b_p, s, x, b_{p+2}, b_{p+1}\}$	net $\{b_{p-1}, b_p, s, x, b_q, b_{q-1}\}$	†-AT $T(s : x : b_{p-1}, b_p \ldots b_{q-1}, b_q)$

Table II. $T(s : c_1, c_2 : l, r)$ is a ‡-AT.

	$q = p+1$	$q = p+2$	$q > p+2$
$p = 0$	4-hole (xc_2b_1lx)	tent $\{l, x, s, c_2, b_2, b_1\}$	‡-AT $T(s : x, c_2 : l, b_1 \ldots b_{q-1}, b_q)^2$
$p = 1$	whipping top $\{l, b_1, x, s, c_2, b_3, b_2\}^1$	net $\{l, b_1, s, x, b_3, b_2\}$	†-AT $T(s : x : l, b_1 \ldots b_{q-1}, b_q)^2$
$p > 1$	long-claw $\{b_{p-2}, b_{p-1}, b_p, s, x, b_{p+2}, b_{p+1}\}$	net $\{b_{p-1}, b_p, s, x, b_q, b_{q-1}\}$	†-AT $T(s : x : b_{p-1}, b_p \ldots b_{q-1}, b_q)$

1 : a 4-hole $(xb_pb_{p+1}b_{p+2}x)$ will be introduced if $x \sim b_{p+2}$;
2 : x is in the category "full" if $q = d+1$.

Table 1: Structures used in the proof of Lemma 5.1 (category "partial")

As the graph is prereduced and contains no small forbidden induced subgraph, it is immediate from Table 1 that the case $q > p+2$ holds, otherwise there always exists a small forbidden induced subgraph (see Table 1). This completes the categorization of vertices of $N(s) \setminus T$. ☐

---

[4] We omit the figure for ‡-ATs: For a ‡-AT $(s : c_1, c_2 : l, r)$, we are only concerned with the relation between center c_2 and $B \cup \{l\}$, and this is the same as the relation between c and $B \cup \{l\}$ in a †-AT.

The proof of our main result of this section is an inductive application of Lemma 5.1. To avoid the repetition of the essentially same argument for †-ATs and ‡-ATs, especially for the interaction between ATs, we use a generalized notation to denote both. We will uniformly use c_1, c_2 to denote center(s) of an AT, and while the AT under discussion is a †-AT, both c_1 and c_2 refer to the only center c. As long as the adjacency of c_1 and l, c_2 and r, and c_1 and c_2 will not concern us here, this unified (abused) notations will not introduce inconsistencies.

Theorem 5.2. *Let* T *be a †- or ‡-AT in a prereduced graph* G *with shallow terminal* s *and base* B*. Let* $C = N(s) \cap N(B)$ *and let* M *be the connected component of* $G - C$ *containing* s*. Then* M *is completely connected to* C*, and* $G[C]$ *is a clique.*

Proof. Denote by $T = (s : c_1, c_2 : l, B, r)$, where $c_1 = c_2 = c$ when T is a †-AT. We show $x \sim y$ for any pair of vertices $x \in C$ and $y \in M$. By definition, $G[M]$ is connected, and there is a chordless path $P = (v_0 \dots v_p)$ from $v_0 = s$ to $v_p = y$ in $G[M]$. We claim that $P \not\sim B$. Suppose the contrary and let v_q be the first vertex in P that is adjacent to B. This means that every v_i with $i < q$ is in category "none" of Lemma 5.1(2). Therefore, applying Lemma 5.1(1,2) on v_i and AT $(v_{i-1} : c_1, c_2 : l, B, r)$ inductively for $i = 1, \dots, q-1$, we conclude that there is an AT $T_i = (v_i : c_1, c_2 : l, B, r)$ for each $i < q$. One more application of Lemma 5.1(1) shows that v_q is adjacent to the center(s) of T_{q-1} as well. If v_q is adjacent to all vertices of B (the category "full" with respect to every T_i), then Lemma 5.1(2) on v_q and T_{q-1} implies that v_q is adjacent to $v_{q-2} \in N(v_{q-1})$, contradicting the assumption that P is chordless. Otherwise (the category "partial"), according to Lemma 5.1(2), there is another AT $T' = (v_{q-1} : c'_1, c'_2 : l', B', r')$, where $B' \subset B$, and $v_q \in \{c'_1, c'_2\}$. Now an application of Lemma 5.1(1) on v_q and T' shows that v_q is adjacent to $v_{q-2} \in N(v_{q-1})$, again a contradiction.

Therefore, $P \not\sim B$ and every v_i is in the category "none" with respect to T. Applying Lemma 5.1 inductively as in the previous paragraph, we get an AT $T_i = (v_i : c_1, c_2 : l, B, r)$ for every $0 \leqslant i \leqslant p$. Vertex x is a neighbor of s and $x \sim B$, thus it cannot be in the category "none." We now separate the discussion based on whether x is in the catergory "full" or "partial." Suppose first that x is in the category "full"; as $x \in N(s)$, Lemma 5.1(1) implies that $x \sim c_1, c_2$. Then applying Lemma 5.1(2) inductively on vertex x, AT T_{i-1}, and vertex $v_i \in N(v_{i-1})$ for $i = 1, \dots, p$ we get that $x \sim v_i$ for each $i \leqslant p$; hence $x \sim v_p = y$. Suppose now that x is in in category "partial." Then by Lemma 5.1(2), there is an AT $T'_0 = (v_0 : c'_1, c'_2 : l', B', r')$, where $B' \subset B$, and $x \in \{c'_1, c'_2\}$. As $P \not\sim B$, we have that $v_i \not\sim B'$ for any $0 \leqslant i \leqslant p$, i.e., v_i is in category "none" with respect to T'_0. Therefore, by an inductive application of Lemma 5.1(2) on the vertex v_i and AT $T'_{i-1} = (v_{i-1} : c'_1, c'_2 : l', B', r')$ for $i = 1, \dots, p$, we conclude that there is an AT $T'_p = (v_p : c'_1, c'_2 : l', B', r')$; hence $x \in \{c'_1, c'_2\}$ is adjacent to $v_p = y$.

To show that $G[C]$ is a clique, let x, y be any pair of vertices in C. If x is in category "full" with respect to T, then Lemma 5.1(2) implies that x is adjacent to $y \in N(s)$. Otherwise, if x is in category "partial" with respect to T, then Lemma 5.1(2) implies that there is an AT $T' = (s : c'_1, c'_2 : l', B', r')$ where $B' \subset B$ and $x \in \{c'_1, c'_2\}$. Therefore, by Lemma 5.1(1) on the vertex $y \in N(s)$ and T', we get that $y \sim c_1, c_2$ and hence $y \sim x$. ☐

We remark that the set C is an M-B separator. Now Theorem 2.1 follows from Theorem 5.2: the set M containing s is in a module whose neighborhood is a clique, hence every vertex in M is simplicial.

6 Long holes

This section proves Theorem 2.2 by showing that the holes in a reduced graph are pairwise congenial. During the study of vertices of a hole, their indices become very subtle. To simplify the presentation, we will frequently apply a common technique, that is, to number the vertices of a hole starting from a vertex of special interest for the property at hand. Moreover, indexing two adjacent vertices in a hole will determine the indices of all the vertices in the hole, as well as the ordering used to transverse the hole.

We start from two simple facts on the relations between vertices and holes, from which we derive the relations between two holes, and finally generalize them to multiple holes.

Proposition 6.1. *For any vertex* v *and hole* H *of a prereduced graph* G*, the set of vertices* $N_H[v]$ *are consecutive in* H*. Moreover, either* $N_H[v] = H$ *or* $|N_H[v]| < |H| - 7$*.*

Proof. This assertion vacuously holds true when $v \not\sim H$, $N_H[v] = H$, or $v \in H$ (then $|N_H[v]| = 3$). We show the existence of forbidden induced subgraphs if the statement of the lemma does not hold. We number the vertices of H in a way that $h_0 \sim v$ but $h_1 \not\sim v$, and then find the following three vertices of H, whose existence is clear from our assumption: *a)* h_{p_1} is the first vertex such that $p_1 > 1$ and $h_{p_1} \sim v$; *b)* h_{p_2} is the first vertex such that $p_2 > p_1$ and $h_{p_2} \not\sim v$; and *c)* h_{p_3} is the first vertex such that $p_2 < p_3 < |H|$ and $h_{p_3} \sim v$, or h_0 if $h_i \not\sim v$ for each $i > p_2$, then $p_3 = |H|$. By the selection of p_1, p_2 and p_3, we have $v \sim h_i$ for each $p_1 \leqslant i < p_2$ and $i = 0, p_3$, but $v \not\sim h_j$ for $0 < j < p_1$ or $p_2 \leqslant j < p_3$ (relations between v and h_i where $p_3 < i < |H|$ are immaterial).

Now we examine the distances between the three indices. There is a small hole of length at most 6 ($v h_0 h_1 \dots h_{p_1} v$) if $p_1 < 4$ or ($v h_{p_2 - 1} h_{p_2} \dots h_{p_3} v$) if $p_3 < p_2 + 4$. Thus $4 \leqslant p_1 < p_2 < p_2 + 4 \leqslant p_3$. Nonetheless, there is • a long claw $\{h_0, v, h_{p_1}, h_{p_1-2}, h_{p_1-1}, h_{p_2}, h_{p_2+1}\}$ if $p_2 = p_1 + 1$; • a net $\{h_0, v, h_{p_1-1}, h_{p_1}, h_{p_2+1}, h_{p_2}\}$ if $p_2 = p_1 + 2$; or • a long claw $\{h_1, h_0, v, h_{p_1-1}, h_{p_1}, h_{p_2}, h_{p_2-1}\}$ if $p_2 > p_1 + 2$. None of these forbidden induced subgraphs involves both h_0 and h_{p_3} and thus they exist regardless of $h_{p_3} = h_0$ or not. This contradiction ensures that $N_H[v]$ are consecutive, so are the vertices of $H \setminus N_H[v]$; there is a hole of length at most 10 if $1 \leqslant |H \setminus N_H[v]| \leqslant 7$. ☐

If a vertex v is adjacent to more than 3 but not all consecutive vertices of a hole H, then we can use v as a shortcut to obtain a hole strictly shorter than H. This cannot happen for a shortest hole. Therefore, the following is immediate (recall that we denote by $\widehat{N}(H)$ the common neighbors of H):

Corollary 6.2. *Let* H *be a shortest hole and* $v \notin \widehat{N}(H)$*. Then* $N_H[v] \leqslant 3$.

Note that each hole H in a prereduced graph contains at least 11 vertices. If v is adjacent to all vertices of H, then on any five consecutive vertices of the hole H, we can apply Proposition 4.3. Therefore v is dominating in the closed neighborhood of H.

Corollary 6.3. *Let* H *be a hole in a prereduced graph and* $v \in \widehat{N}(H)$*. Then* v *is adjacent to all vertices in* $N[H] \setminus \{v\}$.

So far we characterized neighbors of holes in a prereduced graph: Any vertex v is adjacent to a (possibly empty) set of consecutive vertices of a hole H; if v is adjacent to all vertices of H, then it is a common neighbor of the closed neighborhood $N[H]$. From these facts we now derive the relations between holes. The following definition and important lemmas attained as such will reveal the whole picture of all the holes in a prereduced graph.

Definition Two holes H_1 and H_2 are called *congenial* (to each other) if each vertex of one hole is a neighbor of the other hole, that is, $H_1 \subseteq N[H_2]$ and $H_2 \subseteq N[H_1]$.

We remark that this definition implies that every hole is congenial to itself. The following observation partially motivates the definition of congenial holes.

Proposition 6.4. *Let* $\mathcal{H}$ *be a set of holes all congenial to* H*, and* $v \in H$*. Then the set of vertices* $N[v]$ *intersects all holes of* $\mathcal{H}$*.*

Since a vertex in a hole cannot be a common neighbor of it, Corollary 6.3 and the definition immediately imply the following properties:

Corollary 6.5. *Let* H_1 *and* H_2 *be two congenial holes in a prereduced graph* G*. Then* $\widehat{N}(H_1) = \widehat{N}(H_2)$*. Moreover, no vertex of* H_1 *(*H_2 *resp.) is a common neighbor of* H_2 *(*H_1 *resp.).*

We analyze next the relation between two holes that are not congenial to each other. It turns out that if not all vertices of a hole H_1 are adjacent to another hole H_2, then the following lemma shows that every vertex of H_1 is adjacent to either all or none of the vertices of H_2.

Lemma 6.6. *Let* H_1 *and* H_2 *be two adjacent holes in a prereduced graph* G *such that* $H_1 \not\subseteq N[H_2]$*. Then each neighbor of* H_2 *in* H_1 *is a common neighbor of* H_2*, that is,* $N_{H_1}[H_2] \subseteq \widehat{N}(H_2)$*. In particular,* H_1 *and* H_2 *are disjoint.*

Proof. Let u be any vertex in $N_{H_1}[H_2]$, which is non-empty by assumption, and let P be the maximal path in H_1 with the property that $u \in P \subseteq N_{H_1}[H_2]$; denote by p the number of vertices of P. Note that H_1 is not contained in $N[H_2]$ by assumption, thus P is indeed a path. Numbering the vertices in H_1 such that $P = u_0 \dots u_{p-1}$ (the ordering of H_1 is immaterial when $p = 1$ and then u_1 can be either neighbor of u_0 in H_1), the selection of P means $u_i \sim H_2$ for each $0 \leqslant i < p$, and $u_{-1}, u_p \not\sim H_2$ (it is immaterial whether $u_{-1} = u_p$ or not). In the following, we show both ends of P belong to $\widehat{N}(H_2)$, which induces a clique (Proposition 4.5). Thus either $u_0 = u_{p-1}$ (i.e., $p = 1$) or u_0 and u_{p-1} are adjacent (i.e., $p = 2$); in either case, we have $u \in \{u_0, u_{p-1}\} \subseteq \widehat{N}(H_2)$. This proves the first assertion, and the second assertion ensues, as otherwise their common vertices will be common neighbors of H_2, which is not possible.

Note that $u_0 \notin H_2$, as otherwise both neighbors of u_0 in H_1 are adjacent to H_2, contradicting the maximality of P. Similarly, $u_{-1}, u_{-2} \notin H_2$. If u_0 has a unique neighbor v in H_2, then the subgraph induced by u_{-1}, u_0 and five consecutive H_2 vertices centered at v is a long claw (see Figure 4). Now we consider the case $2 \leqslant |N_{H_2}[u_0]| \leqslant |H_2| - 7$ (Proposition 6.1), and number the vertices of H_2 such that $N_{H_2}[u_0] = \{v_1, v_2 \dots, v_q\}$. If u_{-2} is adjacent to v_0, v_1, v_q, or v_{q+1}, then there is a hole $(u_{-2}u_{-1}u_0v_1v_0u_{-2})$, $(u_{-2}u_{-1}u_0v_1u_{-2})$, $(u_{-2}u_{-1}u_0v_qu_{-2})$, or $(u_{-2}u_{-1}u_0v_qv_{q+1}u_{-2})$, respectively. Otherwise, if u_{-2} is not adjacent to $\{v_0, v_1, v_q, v_{q+1}\}$, then there is a long claw $\{u_{-2}, u_{-1}, u_0, v_0, v_1, v_{q+1}, v_q\}$ (see Figure 4, vertex u_p). This proves $u_0 \in \widehat{N}(H_2)$, and with a symmetrical argument we can also prove $u_{p-1} \in \widehat{N}(H_2)$. ☐



Figure 4: u_i ($0 \leqslant i \leqslant p$) has to be a common neighbor of H_2 ($u_{-1}, u_{p+1} \not\sim H_2$)

We are now ready to prove that the congenial relation is transitive. The reflexivity and symmetry of this relation is clear from definition; therefore congenial holes form an equivalent class.

Lemma 6.7. *Let H, H_1 and H_2 be three holes in a prereduced graph G. If both H_1 and H_2 are congenial to H, then H_1 and H_2 are congenial.*

Proof. If H_1 and H_2 are adjacent but not congenial, then Lemma 6.6 implies that one of the holes contains a common neighbor of the other hole. By definition of congenial holes and applying Corollary 6.3 twice, a common neighbor of H_2 is also a common neighbor of H, and also a common neighbor of H_1, and vice versa. Therefore, $H_1 \sim H_2$ leads to a contradiction, as a hole cannot contain a common neighbor of itself.

Assume therefore that there is no edge between H_1 and H_2. Take any vertex $h \in H$. By definition of congenial holes and according to Corollary 6.5, h is adjacent to some but not all vertices of both H_1 and H_2. Now we number the vertices of H_1 and H_2 such that $N_{H_1}[h] = \{u_1, \dots, u_p\}$ and $N_{H_2}[h] = \{v_1, \dots, v_q\}$. There is • a long-claw $\{v_1, h, u_{-1}, u_0, u_1, u_2, u_3\}$ if $p = 1$; • a net $\{v_1, h, u_0, u_1, u_3, u_2\}$ if $p = 2$; or • a long-claw $\{v_0, v_1, u_0, u_1, h, u_p, u_{p+1}\}$ if $p \geqslant 3$. ☐

To prove Theorem 2.2, we show that if there are two holes that are not congenial, then one of them is contained in a nontrivial module. This is impossible in a reduced graph, where every nontrivial module induces a clique. We construct this nontrivial module with the help of the following lemma, which shows that the common neighbors form a separator.

Lemma 6.8. *Let H be a hole that is the shortest among all the holes congenial to it in a prereduced graph G. Then the set of common neighbors $\widehat{N}(H)$ of H separate $N[H] \setminus \widehat{N}(H)$ and $V \setminus N[H]$.*

Proof. Assume to the contrary, $N[H] \setminus \widehat{N}(H)$ and $V \setminus N[H]$ are still connected in $G - \widehat{N}(H)$, then there are two adjacent vertices u and v such that $u \in N[H] \setminus \widehat{N}(H)$ and $v \in V \setminus N[H]$. Note that $u \notin H$, and we have two adjacent vertices only one of which is adjacent to part of the hole H. Depending on the number of neighbors of u in H, we have either a long claw (when $|N_H(u)| = 1$), a net (when $|N_H(u)| = 2$), or a †-AT whose base has five vertices (when $|N_H(u)| = 3$). They cannot exist in a prereduced graph, and on the other hand, if $|N_H(u)| > 3$ then H cannot be the shortest hole by Corollary 6.2. ☐

We are now ready to prove Theorem 2.2:

Proof (of Theorem 2.2). Let H_1 and H_2 be two holes that are not congenial to each other. By Lemma 6.7, being congenial is an equivalence relation. For $i = 1, 2$, let $\mathcal{H}_i$ be the equivalence class where H_i belongs to and let H'_i be a shortest hole in this class. Clearly, H'_1 and H'_2 are not congenial; assume without loss of generality that H'_2 has a vertex v not in $N[H'_1]$. It is clear that H'_1 is shortest among all holes it is congenial to, thus Lemma 6.8 implies that $\widehat{N}(H'_1)$ separates $N[H'_1] \setminus \widehat{N}(H'_1)$ and $V(G) \setminus N[H'_1]$. As $v \notin N[H'_1]$, the set $N[H'_1] \setminus \widehat{N}(H_1)$ is a nontrivial module that does not induce a clique. Thus Reduction 2 is applicable and the graph is not reduced. ☐

7 Breaking holes

A set of vertices is a *hole cover* if it intersects each hole in the graph, and the removal of any hole cover makes the graph chordal. A hole cover is minimal if any proper subset of it is not a hole cover. Any interval deletion set makes a hole cover of the input graph, and thus contains a minimal hole cover. The goal of this section is to prove Theorem 2.3, that is, to provide a polynomial bound on the number of minimal hole covers in a reduced graph and give a polynomial time algorithm to find all of them.[5]

Clearly, a minimal hole cover does not contain any vertex that is not on any hole; this observation will simplify the task of finding all minimal hole covers in a reduced graph.

Proposition 7.1. *Let $\mathcal{H}$ be the set of all holes in a reduced graph G, and G_0 be the subgraph induced by $\bigcup_{H \in \mathcal{H}} H$. A set HC of vertices is a minimal hole cover of G if and only if it is a minimal hole cover of G_0.*

In this section we will focus on the subgraph G_0 induced by the union of all holes in the reduced graph G. The subgraph G_0 has the same set of holes as G, and they remain congenial. Moreover, each vertex of G_0 is in the closed neighborhood of each hole H of G_0.

Proposition 7.2. *The subgraph $G_0 - HC$ is an interval graph for each hole cover HC.*

Proof. Each vertex of G_0 belongs to some hole, and thus cannot be simplicial. Therefore, by Theorem 2.1, G_0 contains no AT. By definition, $G_0 - HC$ contains no hole, thus $G_0 - HC$ is an interval graph. ☐

In particular, by Proposition 6.4, the subgraph $G_0 - N[v]$ is an interval graph for each vertex v of G_0.

We prove a series of claims on how the neighborhood of a vertex v of a hole H_1 looks like in some other hole H_2. The following statement is a paraphrase of Corollary 6.5:

Corollary 7.3. *No vertex v of G_0 can be a common neighbor of any hole in G.*

Therefore, by definition of congenial holes and Proposition 6.1, we can assume that for every $v \in V(G_0)$ and hole H, we have that $N_H[v]$ is a proper nonempty subset of H and its vertices induce a path in H. Fixing any ordering of the vertices in H, we can denote two ends of the path as $\texttt{begin}_H(v)$ and $\texttt{end}_H(v)$ respectively (when $N_H[v]$ contains both h_0 and $h_{|H|-1}$, we number vertices of $N_H[v]$ as $\{h_{-p}, \ldots, h_0, \ldots, h_q\}$ where both p and q are non-negative, and then $\texttt{begin}_H(v) = -p$, $\texttt{end}_H(v) = q$).

Lemma 7.4. *Let H_1 and H_2 be two holes in a reduced graph, and u_0, u_1 be any two adjacent vertices of H_1. The sets of neighbors of u_0 and u_1 in H_2 have the following properties:*

(1) $N_{H_2}[u_0] \cap N_{H_2}[u_1] \neq \emptyset$*;*

(2) *if* $N_{H_2}[u_1] \subseteq N_{H_2}[u_0]$ *or* $N_{H_2}[u_0] \subseteq N_{H_2}[u_1]$*, then either* $\texttt{begin}_{H_2}(u_0) = \texttt{begin}_{H_2}(u_1)$ *or* $\texttt{end}_{H_2}(u_0) = \texttt{end}_{H_2}(u_1)$*; and*

(3) $N_{H_2}[u_0] \cup N_{H_2}[u_1] \neq H_2$.

---

[5] Very recently, Grippo and Safe, in a unpublished manuscript, announce a characterization of forbidden induced subgraphs for normal Helly circular-arc graphs [13, Theorem 6]. It is a simple task to verify that the subgraphs G_0 induced by the union of holes in a reduced graph does not contain any of them. Therefore, there are no three arcs that cover the circle model, and each minimal hole cover corresponds to a point in the circular-arc model. This yields a linear upper bound on the number of minimal hole covers for subgraph G_0 as well as a linear time algorithm to find them.

Proof. According to Proposition 6.1, neighbors of u_0 and u_1 in H_2 are both consecutive. We number the vertices of H_2 such that $N_{H_2}[u_0] = \{v_0, \dots, v_{\ell_1}\}$ and $N_{H_2}[u_1] = \{v_{\ell_2}, \dots, v_{\ell_3}\}$.

On the first assertion, the neighbors of u_0, u_1 in H_2 overlap when either or both of them also belong to H_2, and thus we assume otherwise. Suppose the contrary and $N_{H_2}[u_0] \cap N_{H_2}[u_1] = \emptyset$, then $\ell_1 < \ell_2 \leqslant \ell_3 < |H_2|$. However, either $\ell_3 \geqslant |H_2| - 3$ and $(u_0 u_1 v_{\ell_3} \dots v_{|H_2|} u_0)$ is a hole of length at most 6, or $H = (u_0 u_1 v_{\ell_2} \dots v_{\ell_1} u_0)$ is a hole not congenial to H_2 (in particular, the H_2 vertex $v_{|H_2|-2} \notin N[H]$). This contradicts Theorem 2.2, and thus the first assertion must hold true.

On the second assertion, suppose the contrary and $0 < \ell_2 \leqslant \ell_3 < \ell_1$. If $u_0 \in H_2$ then $\ell_1 = 2$ and $\ell_2 = \ell_3 = 1$; however, applying the first assertion on u_1 and u_2 we can conclude $u_0 (= v_1) \sim u_2$, which is impossible. If $u_1 \in H_2$ then $\ell_1 > \ell_3 \geqslant 3$ and thus $u_0 \notin H_2$; the other neighbor $u_2 \notin H_2$ as otherwise by assumption, $u_0 \sim u_2$. The small minimal forbidden induced subgraphs constructed in the following will hold independent of whether $u_1 \in H_2$ or not. If $u_2 \sim v_0$ then $(v_0 u_0 u_1 u_2 v_0)$ is a 4-hole; if $u_2 \sim v_{-1}$ then $(v_0 u_0 u_1 u_2 v_{-1} v_0)$ is a 5-hole. Thus we conclude $u_2 \not\sim v_0, v_{-1}$, and with the same argument we conclude $u_2 \not\sim v_{\ell_1+1}, v_{\ell_1}$. Nonetheless $\{u_2, u_1, u_0, v_{\ell_1+1}, v_{\ell_1}, v_{-1}, v_0\}$ will be a long-claw: $(v_{\ell_1+1} v_{\ell_1} u_0 v_0 v_{-1})$ is clearly a chordless path, and u_1 is only adjacent to u_0 in it. Therefore, either $0 = \mathtt{begin}_{H_2}(u_0) = \mathtt{begin}_{H_2}(u_1) = \ell_2$ or $\ell_3 = \mathtt{end}_{H_2}(u_0) = \mathtt{end}_{H_2}(u_1) = \ell_1$, and this proves the second assertion.

Suppose the third assertion is not true and the neighbors of u_0 and u_1 in H_2 cover the entire hole H_2. According to Proposition 6.1 and Corollary 6.5, there are at least 7 vertices in both $N_{H_2}[u_0] \setminus N_{H_2}[u_1]$ and $N_{H_2}[u_1] \setminus N_{H_2}[u_0]$; this implies $u_0, u_1 \notin H_2$. Then $\{v_{\ell_1}, v_{\ell_1+1}, v_{\ell_1+2}, \dots, v_{|H_2|-1}, v_{|H_2|}\} \subseteq N_{H_2}[u_1]$: there is a 4-hole $(u_0 v_{\ell_1} v_{\ell_1+1} u_1 u_0)$ or $(u_0 v_{|H_2|} v_{|H_2|-1} u_1 u_0)$ if $u_1 \not\sim v_{\ell_1}$ or $u_1 \not\sim v_{|H_2|} = v_0$. According to Proposition 6.1, $\ell_1 \geqslant \ell_2 > 6$ and $\ell_3 - \ell_2 \geqslant \ell_3 - \ell_1 > 6$. Let $2 < p < |H_1| - 1$ be the index of the first H_1 vertex that is either adjacent to $N_{H_2}[u_0]$ but neither v_0 nor v_{ℓ_1}, or adjacent to $N_{H_2}[u_1]$ but neither v_{ℓ_2} nor v_{ℓ_3}.

To show the existence of p, we start from examining the neighbors of u_3 in H_2, which has to be a subset of $N_{H_2}[u_1]$, otherwise there is a 4-hole $(v_i u_0 u_1 u_2 v_i)$ or 5-hole $(v_i u_0 u_1 u_2 u_3 v_i)$, depending on $u_2 \sim v_i$ or not, if $u_3 \sim v_i$ for some $v_i \notin N_{H_2}(u_1)$. As $u_1 \not\sim u_3$ and then $|N_{H_2}[u_3]| \leqslant 2$ (Corollary 6.3). If $p \neq 3$, then u_3 is adjacent one end of $N_{H_2}[u_1]$, and without loss of generality, let it be v_{ℓ_2}. By assumption, v_{ℓ_2} is also in $N_{H_2}[u_0]$, thus it can be inferred from $p \neq 3$ that u_3 is adjacent to one end of $N_{H_2}[u_0]$, which has to be v_{ℓ_1} and $\ell_1 = \ell_2$ or $\ell_1 = \ell_2 + 1$ (we are using the facts $\ell_1 \geqslant \ell_2 > 6$ and $|N_{H_2}[u_3]| \leqslant 2$). According to Corollary 6.5, neither v_{ℓ_1} nor v_{ℓ_2} can be a common neighbor of H_1, which implies that there must be a vertex u_p for $2 < p < |H_2| - 1$ that is not adjacent to one of v_{ℓ_1} and v_{ℓ_2}, and we claim that vertex satisfies the condition. As $u_{p-1} \not\sim u_0, u_1$, it is adjacent to neither v_{ℓ_1-2} nor v_{ℓ_2+2}. and $N_{H_2}(u_{p-1}) \subseteq \{v_{\ell_1-1}, v_{\ell_1}, v_{\ell_2}, v_{\ell_2+1}\}$. By the first assertion and the fact that u_p is not adjacent to both v_{ℓ_1} and v_{ℓ_2}, either $N_{H_2}[u_p] \subset N_{H_2}[u_0]$ or $N_{H_2}[u_p] \subset N_{H_2}[u_1]$. Therefore $|N_{H_2}[u_p]| \leqslant 2$ (as $u_p \not\sim u_0, u_1$), and $u_p \not\sim v_0$ (as $\ell_1 > 6$).

Without loss of generality, assume u_p to be adjacent to $N_{H_2}[u_1]$ but neither v_{ℓ_2} nor v_{ℓ_3}. By Proposition 4.3 and the fact $u_p \not\sim u_1$, we conclude $u \not\sim v_i$ for any $\ell_2 + 2 \leqslant i \leqslant \ell_3 - 2$. As a result, $N_{H_2}[u_p] = \{v_{\ell_2+1}\}$ or $N_{H_2}[u_p] = \{v_{\ell_3-1}\}$. There is a net $\{v_{\ell_2-1}, v_{\ell_2}, u_p, v_{\ell_2+1}, v_{\ell_2+3}, u_1\}$ of $\{v_{\ell_3+1}, v_{\ell_3}, u_p, v_{\ell_2-1}, v_{\ell_2-3}, u_1\}$ (noting $\ell_3 - \ell_2 > 6$). This is impossible, and thus the third assertion must hold true. ☐

This implies the following corollaries.

Corollary 7.5. *Let* H_1 *and* H_2 *be two holes in a reduced graph, and* u_0, u_1 *be any two adjacent vertices of* H_1. *Then* u_1 *is adjacent to at least one of* $\mathtt{begin}_{H_2}(u_0)$ *and* $\mathtt{end}_{H_2}(u_0)$.

Corollary 7.6. *Let* H_1 *and* H_2 *be two holes in a reduced graph, and* P *be a path in* H_1. *Then* $\bigcup_{u \in P} N_{H_2}[u]$ *are consecutive.*

Lemma 7.7. *Any minimal hole cover of a reduced graph* G *induces a clique.*

Proof. Assume to the contrary, there is a minimal hole cover HC that contains two non-adjacent vertices u and v. By the minimality of HC, there are two holes H_u and H_v in G such that $HC \cap H_u = \{u\}$ and $HC \cap H_v = \{v\}$. Note that $u \notin H_v$ and $v \notin H_u$. Let us number the vertices of H_u such that $\mathtt{begin}_{H_u}(v) = 0$ and $\mathtt{end}_{H_u}(v) = p$, where $p = |N_{H_u}(v)| - 1$. As both $N_{H_v}[u_0]$ and $N_{H_v}[u_p]$ contain v and consecutive in H_v by Proposition 6.1, their union $N_{H_v}[u_0] \cup N_{H_v}[u_p]$ is also consecutive in H_v. We number the vertices of H_v such that $u_0 \sim v_0$ and $N_{H_v}[u_0] \cup N_{H_v}[u_p] = \{v_0, v_1, \dots, v_q\}$, where $q = |N_{H_v}[u_0] \cup N_{H_v}[u_p]| - 1$. We argue that $q < |H_v| - 1$, i.e. $N_{H_v}[u_0] \cup N_{H_v}[u_p]$ cannot cover H_v. This follows from Lemma 7.4(3) when $p = 1$. Now suppose $p > 1$, which implies $u_0 \not\sim u_p$, and $q = |H_v| - 1$. Recall that by Proposition 6.1, $7 < |N_{H_v}[u_0]| < |H_v| - 7$. We consider the two ends $\mathtt{begin}_{H_v}[u_0]$ and $\mathtt{end}_{H_v}[u_0]$, one of which must have distance more than 2 to v in H_v. Without loss of generality, let it be $v_i = \mathtt{end}_{H_v}[u_0]$. Then either v_i is



Figure 5: There is a hole avoiding both u and v: it cannot contain chord $(u_j v_i)$.

also adjacent to u_p, then there is a 4-hole $(v u_0 v_i u_p v)$; or $v_i \not\sim u_p$ and $v_{i+1} \sim u_p$, then there is a 5-hole $(v u_0 v_i v_{i+1} u_p v)$.

Denote by $P_u = (u_0 u_1 \dots u_p)$ and $P_v = (v_q v_{q+1} \dots v_0)$. By definition $u \notin P_u$. Likewise, $v \notin P_v$: by definition, $v \in \{v_0, v_1, \dots, v_q\}$, but v cannot be v_0 or v_q, since Corollary 7.5 implies that the two neighbors of v in H_v are adjacent to either u_0 or u_p (and v_{-1} and v_{q+1} are not in $N_{h_v}[u_0] \cup N_{h_v}[u_q]$ by definition). In what follows we show the existence of a hole in $G - HC$, which contradicts the fact that HC is a hole cover.

We first show $v_i \not\sim P_u$ for each $i > q$ by contradiction (see red edges in Figure 5). Suppose the contrary, if v_i is adjacent to a vertex of P_u, then the facts $v_i \not\sim u_0$, $v_i \not\sim u_p$ and Proposition 6.1 imply that every neighbor of v_i in H_u is in $\{u_1, \dots, u_{p-1}\}$. Note that $v_i \notin H_u$, as otherwise it must belong to P_u and thus adjacent to v, which is impossible. We argue that $|N_{H_u}[v_i]| = 1$ and the only neighbor has to be u_1 or u_{p-1}: as $v_i \not\sim v$, it follows from Proposition 4.4. There is • a †-AT $(v_i : u_1 : u_{-1}, u_0 v u_2, u_3)$ with size 7 if $p = 2$; or • a net $\{u_{-1}, u_0, v_i, u_1, u_3, v\}$ or $\{u_{p+1}, u_p, v_i, u_{p-1}, u_{p-2}, v\}$ if $p > 2$. This is impossible, hence $v_i \not\sim P_u$ for each $i > q$.

Now we construct the hole in $G - HC$ as follows (see red edges in Figure 5). Recall that the length of P_v is at least 2. If $u_0 \sim v_q$ then $(u_0 P_v u_0)$ is such a hole. Otherwise by assumption we have $u_p \sim v_q$. Let $\ell_1 = \max\{i | u_i \sim v_0 \text{ and } 0 \leqslant i \leqslant p\}$, and $\ell_2 = \min\{i | u_i \sim v_q \text{ and } \ell_1 \leqslant i \leqslant p\}$. Then $(u_{\ell_1} u_{\ell_2} P_v u_{\ell_1})$ will be such a hole. □

As a consequence, any minimal hole cover of a reduced graph G_0 intersects each hole in it by either one or two vertices.

Lemma 7.8. *Let* HC *be a minimal hole cover of a reduced graph* G*. There is a vertex* v *in* G_0 *such that* $N_{G_0}[v] \not\sim HC$.

Proof. We show this by construction. Let H be a shortest hole in $\mathcal{H}$. By Lemma 7.7, there are at most two vertices of $H \cap HC$, and they are consecutive in H. We number the vertices of H such that $h_0 \in HC$ and $h_1 \notin HC$. We claim $v = h_5$ is the asserted special vertex. Recall that by Corollaries 6.2 and 7.3, each vertex of G_0 has at most 3 neighbors on H. Assume to the contrary, $N_{G_0}[h_5]$ and the clique HC (Lemma 7.7) are adjacent, then there is an h_0-h_5 path P of length at most 3 and all its interior vertices belong to G_0. The case $P = h_0 v h_5$ is impossible, as by Proposition 6.1 and Corollary 6.2, v is adjacent to at most 3 consecutive vertices in H. Now we may assume $P = h_0 v_1 v_2 h_5$, and examine the neighbors of v_1 and v_2 in H. Let h_i be the last neighbor of v_1 in H, and h_j be the first neighbor of v_2 in H. By Corollary 6.2, we have $i \leqslant 2$ and $j \geqslant 3$. Therefore there is a hole $(v_1 h_i h_{i+1} \dots h_j v_2 v_1)$ of length at least 4 and at most 8. □

We prove Theorem 2.3 by showing that we can enumerate hole covers by enumerating separators in an interval graph. In one direction of the proof, we need the following claim. Recall that a connected interval graph corresponds to a consecutive set of intervals in the real line, and a point p in the real line that has intervals not containing p and lying at its both sides decides a clique separator, that is, the set of vertices whose corresponding intervals contain p.

Proposition 7.9. *Let v be a vertex in an interval graph G. If v is not adjacent to any simplicial vertex, then $N[v]$ is a separator of G.*

Proof. We consider the interval representation of G. Without loss of generality, we assume that no two intervals have the same ends. Denote by x the interval with the smallest right end, and y the interval with the largest left end. It is easy to see that x and y are simplicial. Furthermore, if $x \sim y$, then the graph is a complete graph (every interval contains the interval between the left end of y and the right end of x), and every vertex is adjacent to a simplicial vertex, thus the statement is vacously true. Therefore, we can assume $x \not\sim y$. For any point p in the interval corresponding to v, the set $N[v]$ contains all vertices whose intervals contain p. Neither x nor y is adjacent to v, and then $N[v]$ is an x-y separator. ☐

The following lemma formulates the equivalence of hole covers and separators in our setting:

Lemma 7.10. *Let G be a reduced graph, and G_0 be the subgraph induced by all holes of G. Let v be a vertex of G_0, and X be a set of vertices of G_0 that are not adjacent to $N_{G_0}[v]$. Then X forms a minimal hole cover of G if and only if it is a minimal separator of $G_0 - N[v]$.*

Proof. According to Proposition 7.1, it suffices to show that X is a hole cover of G_0 if and only if it is a separator of $G_0 - N[v]$ (which is the same as $G_0 - N_{G_0}[v]$). According to Propositions 6.4 and 7.2, $G_0 - N[v]$ is an interval graph.

$\Rightarrow$ The set X is a trivial separator of $G_0 - N[v]$ if it is not connected: there are at least two connected components in $G_0 - N[v]$, each of which contains neighbors of $N[v]$ and thus the number of connected components of $G_0 - N[v] - X$ is no less than that of $G_0 - N[v]$. Hence in the argument of this direction we assume $G_0 - N[v]$ is connected, and an alternative way of phrasing that “the hole cover X is a separator of the connected subgraph $G_0 - N[v]$” is to say that $G_0 - (N[v] \cup X)$ is not connected, or $N[v]$ is a separator of $G_0 - X$. By Proposition 7.2, $G_0 - X$ is an interval graph as well. By definition, there is no simplicial vertex in G_0; hence each simplicial vertex x in $G_0 - X$ is adjacent to X, as otherwise $N_{G_0 - X}(x) = N_{G_0}(x)$ and is not a clique. As $N[v]$ is not adjacent to X by assumption, v is not adjacent to any simplicial vertex. Therefore, according to Proposition 7.9, the removal of $N[v]$ disconnects the interval graph $G_0 - X$, and the “only if ” direction is proved.

$\Leftarrow$ Suppose the contrary and there is a hole H in $G_0 - X$. By the construction of G_0, the vertex v is contained in some hole and this hole is congenial to H (G is a reduced graph); then by definition of congenial holes and Corollary 6.5, v is adjacent to some but not all vertices of H. We number the vertices of H such that $h_0 \in N[v]$ and $h_1 \notin N[v]$, and denote by P the path $H \setminus N_H[v]$. As H is not destroyed by X, that is, $P \cap X = \emptyset$; the vertices of P are contained in the same component C in the subgraph $G_0 - N[v] - X$. We show that each vertex $u \in G_0 - N[v] - X$ is also in C, contradicting the assumption that X is a separator. Let H_2 be any hole that contains u. Vertices of $H_2 \setminus X$ adjacent to P are clearly in the component C. Since H and H_2 are congenial, each vertex of H_2 not adjacent to P has to be adjacent to $N_H[v]$. Let $U \subseteq H_2$ be the set of vertices adjacent to $N_H[v]$. According to Corollary 7.6, set U induces a path P_2. As $X \not\sim N[v]$ by assumption, the path P_2 is disjoint from X. Thus we can finish the proof by showing that P_2 is also in C. According to Lemma 7.4(1), $N_{H_2}[h_0] \cap N_{H_2}[h_1] \neq \emptyset$, i.e. there is a vertex $u' \in H_2$ that is adjacent to both h_0 and h_1. Clearly, vertex u' is in P_2 (being a neighbor of h_0) and in component C (being a neighbor of h_1). Therefore, the subgraph $G_0 - N[x] - X$ contains only one component, contradicting the fact that X is a separator of $G_0 - N[x]$. ☐

We are now ready to prove Theorem 2.3:

Proof (of Theorem 2.3). Let G_0 be induced by the union of the holes of G. On one hand, according to Lemmas 7.8 and 7.10, each minimal hole cover of G corresponds to a minimal separator of $G_0 - N[v]$ for some vertex v of G_0. On the other hand, there are at most n minimal separators in $G_0 - N[v]$ for each vertex v of G_0, hence a quadratic bound for the total number of minimal hole covers of G. To enumerate all these hole covers, we try every vertex v of G_0, compute an interval representation of the interval graph $G_0 - N[v]$, and enumerate all its minimal separators. ☐

8 Caterpillar decompositions

We prove Theorem 2.4 in this section by providing an FPT algorithm for INTERVAL DELETION under the assumption that the graph is prereduced, chordal, and every shallow terminal is simplicial. The algorithm presented in this section finds an AT satisfying a certain minimality condition, and we show that it contains a set of 10 vertices such that there is a minimum interval deletion set containing at least one of these 10 vertices. Therefore, we can branch on deleting one of these 10 vertices. Note that after any vertex deletion, the graph remains chordal, prereduced, and every shallow terminal is still simplicial, as all these properties are hereditary.

Denote by $\mathsf{ST}(G)$ the set of shallow terminals, which are a subset of simplicial vertices. Using the properties of simplicial vertices and according to Proposition 4.2, it is easy to check that for each †- or ‡-AT, its shallow terminal is in $\mathsf{ST}(G)$, its base terminals might or might not be in $\mathsf{ST}(G)$ (see Figure 9 for examples), and all non-terminal vertices cannot be in $\mathsf{ST}(G)$. The following proposition can be derived from Lemma 5.1. Indeed, starting from any AT T with shallow terminal s we can find such an AT: no vertex of $N(s) \setminus \mathsf{ST}(G)$ can be in the "none" category w.r.t. any AT; as long as there is another vertex $v \in N(s) \setminus \mathsf{ST}(G)$ in the "partial" category, we can find another AT with a base $B' \subset B$.

Proposition 8.1. *Let G be a prereduced chordal graph where all shallow terminals are simplicial, and $s \in \mathsf{ST}(G)$. There is an AT whose base B are adjacent to all vertices of $N(s) \setminus \mathsf{ST}(G)$.*

Now that the graph is chordal, it makes sense to discuss its clique tree, which shall be the main structure of this section. Since the interior vertices of any shortest path cannot be simplicial, the removal of simplicial vertices will not disconnect a connected graph; as long as G is connected, $G - \mathsf{ST}(G)$ will be a connected interval graph. This observation suggests that the clique tree of G has a very nice structure. A *caterpillar (tree)* is a special tree that consists of a central path and all other vertices are leaves connected to it.

Proposition 8.2. *Let G be a prereduced chordal graph where all shallow terminals are simplicial. There is a clique tree $\mathcal{T}(G)$ that is a caterpillar, where all shallow terminals appear only in leaf nodes, and each other vertex appears in some node in the central path.*

Proof. Let us inspect every maximal clique K of G. If K contains some vertex $v \in \mathsf{ST}(G)$, then it must be $K = N[v]$: K is a clique, hence $K \subseteq N[v]$; the containment cannot be proper as K is maximal and $N[v]$ induces a clique. If $K \cap \mathsf{ST}(G) = \emptyset$, then K is also a maximal clique of $G - \mathsf{ST}(G)$. On the other hand, a maximal clique K' of $G - \mathsf{ST}(G)$ has to be a maximal clique of G as well; otherwise it is a proper subset of $N[v]$, which, however, according to Proposition 8.1, cannot be maximal in $G - \mathsf{ST}(G)$. Therefore, a maximal clique of G is either $N[v]$ for some vertex $v \in \mathsf{ST}(G)$ or a maximal clique of $G - \mathsf{ST}(G)$. Now we construct the clique tree as follows. We first make a clique path $\mathcal{T}'$ for the interval graph $G - \mathsf{ST}(G)$ (Theorem 3.1), and then attach each maximal clique $N[v]$ of the first kind to $\mathcal{T}'$ at some maximal clique that properly contains $N[v] \setminus \mathsf{ST}(G)$ (arbitrarily pick from multiply choices). Clearly it satisfies the second condition of clique tree. ☐

Fixing a particular clique tree $\mathcal{T}(G)$, we number the nodes (maximal cliques) on the central path as $K_0, K_1, \dots$. By Proposition 8.2 and definition of clique tree, each vertex $v \in V(G) \setminus \mathsf{ST}(G)$ is contained in some consecutive nodes of the central path. Denote by $\mathtt{first}(v)$ and $\mathtt{last}(v)$ the indices of the first and last nodes that contain the vertex $v \notin \mathsf{ST}(G)$. Given any AT T, the vertices of its base B are not simplicial, hence in the main path of the caterpillar decomposition. As the graph is prereduced, B contains at least 6 vertices (note $|T| \geqslant 11$); in particular, the nodes that contain b_1 and b_d are disjoint ($b_1 \not\sim b_d$). Without loss of generality, we assume that $\mathtt{last}(b_1) < \mathtt{first}(b_d)$, i.e., the base B goes from "left to right." In this way base terminals l and r are clearly defined, as $l \sim b_1$ and $r \sim b_d$; so are the centers c_1 and c_2 when T is a ‡-AT as $l \sim c_1$ and $r \sim c_2$. Note that l and r can be in $\mathsf{ST}(G)$, and thus it is possible that $\mathtt{first}(c_1) > \mathtt{first}(c_2)$ and/or $\mathtt{last}(c_1) > \mathtt{last}(c_2)$ when T is a ‡-AT, but this will not matter in the proofs to follow.

The following properties of the clique tree are straightforward by observing the adjacencies and nonadjacencies between the vertices of an AT and the position of the intervals corresponding to vertices not in $\mathsf{ST}(G)$, so we state here without proof. To avoid the repetition of the essentially same argument for †-ATs and ‡-ATs, we follow the same convention as in Section 4.

Proposition 8.3. *Let $T(s : c_1, c_2 : l, B, r)$ be a †- (then $c_1 = c_2 = c$) or ‡-AT. Then*

$$\begin{aligned}\big(\mathtt{first}(b_1) \leqslant{} & \mathtt{last}(b_0) <\big)\ \mathtt{first}(c_2), \mathtt{first}(b_2) \leqslant \mathtt{last}(b_1) \leqslant \dots \\ & \leqslant \mathtt{first}(b_i) \leqslant \mathtt{last}(b_{i-1}) < \mathtt{first}(b_{i+1}) \leqslant \mathtt{last}(b_{i-1}) < \mathtt{first}(b_{i+1}) \\ \leqslant \cdots <{} & \mathtt{first}(b_d) \leqslant \mathtt{last}(b_{d-1}), \mathtt{last}(c_1)\ \big(< \mathtt{first}(b_{d+1}) \leqslant \mathtt{last}(b_d)\big),\end{aligned} \tag{1}$$

where relations in parentheses only hold when $l = b_0 \notin \mathsf{ST}(G)$ and $r = b_{d+1} \notin \mathsf{ST}(G)$ respectively.

The non-terminal vertices of an AT also appear in a consecutive set of nodes in the central path of $\mathcal{T}(G)$; as defined and explained as follows, the minimum set will be of special interest.

Definition Let G be a prereduced chordal graph where every shallow terminal is simplicial and $\mathcal{T}(G)$ be a caterpillar clique tree of G. Given any pair of indices $p \leqslant q$, the (p,q)-*segment* comprises the set of vertices $U_{p,q} = \bigcup_{p \leqslant i \leqslant q} K_i$. Denote by $N_{p,q}$ the set of common vertices $\bigcap_{p \leqslant i \leqslant q} K_i = K_p \cap K_q$. Each AT T of G determines a segment $\mathtt{seg}(T) = U_{\mathtt{last}(b_1),\mathtt{first}(b_d)}$, where b_1 and b_d are ends of the base B of T; such a segment is called a *minimal segment* if there is no AT T' in G such that $\mathtt{seg}(T') \subset \mathtt{seg}(T)$.

By definition of clique trees, $N_{p,q} = K_p \cap K_q$. It is easy to verify that all non-terminal vertices of T are contained in $\mathtt{seg}(T)$, which is the inclusive-wise minimal (also minimum) segment with this property. In particular, $b_1 \notin U_{p,q}$ when $p > \mathtt{last}(b_1)$ and $b_d \notin U_{p,q}$ when $q < \mathtt{first}(b_d)$. Moreover, the set $N_{\mathtt{last}(b_1),\mathtt{first}(b_d)}$ is exactly the set of common neighbors of the base of T. Indeed, by definition $N_{\mathtt{last}(b_1),\mathtt{first}(b_d)} \subseteq \widehat{N}(B)$. If $v \notin N_{\mathtt{last}(b_1),\mathtt{first}(b_d)}$, then either $\mathtt{first}(v) > \mathtt{last}(b_1)$, implying $v \not\sim b_1$, or $\mathtt{last}(v) < \mathtt{last}(b_d)$, implying $v \not\sim b_d$; therefore $\widehat{N}(B) \subseteq N_{\mathtt{last}(b_1),\mathtt{first}(b_d)}$.

Proposition 8.4. *Let* $T = (s : c_1, c_2 : l, B, r)$ *be a* †- *(then* $c_1 = c_2 = c$*) or* ‡*-AT such that* $\mathtt{seg}(T)$ *is a minimal segment. Then* $s \not\sim u$ *for each vertex* $u \in \mathtt{seg}(T) \setminus \widehat{N}(B)$. *Moreover, if* lPr *is a chordless* l-r *path in the subgraph induced by* $\big(\mathtt{seg}(T) \setminus \widehat{N}(B)\big) \cup \{l, r\}$, *then* $T_P = (s : c_1, c_2 : l, P, r)$ *makes an AT, and* $\mathtt{seg}(T_P) = \mathtt{seg}(T)$.

Proof. For the first assertion, suppose the contrary and $s \sim u$, then Lemma 5.1 applies. Observe that u is in the category "partial": it is adjacent to some (because $u \in \mathtt{seg}(T)$) but not all vertices in B (because $u \notin \widehat{N}(B)$). Hence there is another AT T' whose base is a proper subset of B. Note that $\mathtt{last}(b_i) > \mathtt{last}(b_1)$ and $\mathtt{first}(b_i) < \mathtt{first}(b_d)$ for each $1 < i < d$, from which we can conclude $\mathtt{seg}(T') \subset \mathtt{seg}(T)$, a contradiction to the minimality of $\mathtt{seg}(T)$.

To verify that T_P is an AT, it suffices to show that $v \sim c_1$, $v \sim c_2$, but $v \not\sim s$ for each vertex $v \in P$. Observing that $v \in \mathtt{seg}(T) \setminus \widehat{N}(B)$, it is adjacent to center(s) as $c_1, c_2 \in \widehat{N}(B) = N_{\mathtt{last}(b_1),\mathtt{first}(b_d)}$, and not adjacent to s by the first assertion. Since the base P of T_P is fully contained in $\mathtt{seg}(T)$, we conclude $\mathtt{seg}(T_P) \subseteq \mathtt{seg}(T)$; then the last assertion holds true as $\mathtt{seg}(T)$ is minimal. ☐

In Section 6, we considered holes of shortest length and observed that a vertex sees either all or at most 3 consecutive vertices of the hole. Proposition 8.4 allows us to consider ATs whose base is a shortest path, and then we can observe an analogous statement about the number vertices in B a vertex can see.

Definition Let $T = (s : c_1, c_2 : l, B, r)$ be a †- (then $c_1 = c_2 = c$) or ‡-AT such that $\mathtt{seg}(T)$ is a minimal segment. We say B is a *short* base if B is a shortest l-r path in the subgraph induced by $\big(\mathtt{seg}(T) \setminus \widehat{N}(B)\big) \cup \{l, r\}$.

Note that if all non-terminal vertices of an AT are contained in a minimal segment $U_{p,q}$, then its base cannot pass through $N_{p,q}$, otherwise $|B| = 3$. Thus the base B indeed gives a l-r path in the subgraph induced by $\big(\mathtt{seg}(T) \setminus \widehat{N}(B)\big) \cup \{l, r\}$. Therefore, Proposition 8.4 can be used to replace the base with a short base. As this replacement does not change $\mathtt{seg}(T)$ and hence does not break minimality, we can assume from now that we are working with an AT T such that $\mathtt{seg}(T)$ is minimal and T has a short base.

Proposition 8.5. *Let* $T = (s : c_1, c_2 : l, B, r)$ *be a* †- *(then* $c_1 = c_2 = c$*) or* ‡*-AT such that* $\mathtt{seg}(T)$ *is a minimal segment and* B *is a short base. If* $u \in U_{\mathtt{last}(b_3),\mathtt{first}(b_{d-2})} \setminus \widehat{N}(B)$*, then* u *has at most three neighbors in* $B \cup \{l, r\}$*, and there is an index* $3 \leqslant \ell \leqslant d-2$ *such that*

$$\mathtt{first}(c_2) \leqslant \mathtt{last}(b_{\ell-2}) < \mathtt{first}(u) \leqslant \mathtt{last}(u) < \mathtt{first}(b_{\ell+2}) \leqslant \mathtt{last}(c_1). \tag{2}$$

Proof. The neighbors of u on the chordless path lBr have to be consecutive, otherwise there is a hole. If u has more than three neighbors, then we can use u as a shortcut to obtain a strictly shorter l-r path, contradicting the assumption that T has a short base. Vertex u has at least one neighbor in B; let b_p and b_q be the first and last neighbors of u in $B \cup \{b_0, b_{d+1}\}$, respectively. Then selecting $\ell = (p+q)/2$ will justify (2) (observe that $\ell - 2 < p$ and $\ell + 2 > q$, thus $u \not\sim b_{\ell-2}, b_{\ell+2}$. By assumption, $u \sim b_i$ for some $4 \leqslant i \leqslant d-3$, which implies $1 < p < d$, and thus $3 \leqslant \ell \leqslant d-2$. ☐

In the final lemma we will be interested in a special minimal segment that admits one more special property and a representative AT contained in it. Among all minimal segments, the one with the smallest end-index is called the *leftest minimal segment*. This segment is unique as minimal segments cannot properly contain each other. This restriction allows us to consider only the *right side* of this segment: there is no segment defined by another AT that is *left* to it.

Lemma 8.6. *Let* G *be a chordal and prereduced where every shallow terminal is simplicial. Let* $T = (s : c_1, c_2 : b_0, B, b_{d+1})$ *be a* †- *(then* $c_1 = c_2 = c$*) or* ‡*-AT in* G *such that* $\mathtt{seg}(T)$ *is the leftest minimal segment and* B *is a short base. There is a minimum interval deletion set to* G *that either contains one of*

$$V_B = \{s, c_1, c_2, b_0, b_i \mid i \geqslant d-4\},$$

or the set of vertices $X = S_{i,i+1} \setminus \widehat{N}(B)$*, where* $S_{i,i+1}$ *is a minimum separator among* $\mathtt{last}(b_0) \leqslant i < \mathtt{first}(b_{d-4})$.

Proof. Given any minimum interval deletion set Q, we assume $Q \cap V_B = \emptyset$, and $X \not\subseteq Q$; otherwise we have nothing to prove. Denote by $N = \widehat{N}(B)$, and $V_I = \bigcup_{\mathtt{last}(b_1)<i<\mathtt{first}(b_{d-4})} K_i \setminus N$ (note that the bounds on i are strict and b_{d-4} appears in the upper bound). The motivation behind the definition of V_I is that any vertex $v \in V_I$ is in $N[b_i]$ for some $i < d-4$, and any path between v and a vertex right to $\mathtt{seg}(T)$ avoiding N contains at least two vertices of $\mathtt{seg}(T)$. Recall that $c_1, c_2 \in N$. We start from another simple observation.

Claim 1. *For each vertex* $v \in U_{0,\mathtt{first}(b_{d-2})} \setminus N$*, we have* $\mathtt{last}(v) < \mathtt{first}(b_d)$*, and* $v \not\sim ST(G)$.

Proof. We always have $\mathtt{last}(v) < \mathtt{first}(b_d)$: by Proposition 8.5 when $v \in U_{\mathtt{last}(b_3),\mathtt{first}(b_{d-2})} \setminus N$; and by the fact $\mathtt{last}(v) < \mathtt{last}(b_3) < \mathtt{first}(b_d)$ when $v \in U_{0,\mathtt{first}(b_{d-2})} \setminus U_{\mathtt{last}(b_3),\mathtt{first}(b_{d-2})}$. Suppose the contrary and $v \sim x$ for some vertex $x \in ST(G)$. By definition, x is a shallow terminal of some AT T'. Applying Lemma 5.1(2) on v and T', vertex v cannot be in category "none": $v \notin ST(G)$, hence v cannot be a shallow terminal. If v is in categoty "full," then the base of T' is contained in $U_{\mathtt{first}(v),\mathtt{last}(v)}$, contradicting the selection of T. If v is in category "partial," then there is another AT T'' with a v as a center, hence the base of T'' is contained in $U_{\mathtt{first}(v),\mathtt{last}(v)}$, again contradicting the selection of T. ☐

We prove this lemma by constructing a minimum interval deletion set $Q' = (Q \setminus V_I) \cup X$ that contains X. Recall that by definition, the set $X \cup N$ is a separator. As G is already chordal, all forbidden induced subgraphs in all subgraphs are ATs. We now argue this by showing Q' covers all ATs that are covered by Q. Suppose the contrary and there is an AT AT T' that is disjoint from Q', then by the construction of Q', T' must intersect the set $V_I \setminus X$. Let $u \in T' \cap (V_I \setminus X)$. According to Claim 1, u is not adjacent to $ST(G)$ and thus cannot be a center; u is in the base, or one of the base terminals.

Now we consider the base B' of T', which are fully contained in the central path of $\mathcal{T}(G)$, and the segment $\mathtt{seg}(T')$ defined by B'. Recall that center(s) c'_1, c'_2 of T' are dominating in $\mathtt{seg}(T')$. As $\mathtt{seg}(T')$ cannot be left to $\mathtt{seg}(T)$ (i.e. $\mathtt{first}(b'_{d'})$ cannot be smaller than $\mathtt{first}(b_d)$), if $u \in B' \subset \mathtt{seg}(T')$, then we argue that center(s) have to be in N. As a consequence, the base terminal b'_0 has to be left to $\mathtt{seg}(T)$, and the base has to go through X as it cannot go through N. Likewise, we argue B' has to go through N and there exists an AT that avoids Q when $u = b'_0 \notin \mathtt{seg}(T')$. The remainder of this proof breaks into four steps.

Claim 2. $c'_2 \notin N$.

Proof. Suppose the contrary and $c'_2 \in N$, then we show that T' has to intersect X, and thus cannot exist in $G - Q'$. We first show that c'_1 is also in N when T' is a ‡-AT. By the selection of T, the vertex u cannot be the terminal $b'_{d'+1}$, as otherwise according to Proposition 8.3, $\mathtt{first}(b'_{d'}) < \mathtt{first}(u) < \mathtt{first}(b_d)$. Thus $c'_1 \sim u$, which implies $c'_1 \in U_{0,\mathtt{first}(b_{d-2})}$ (Proposition 8.5), and then $c'_1 \in N$ (Claim 1). Now that the center(s) are in N and accordingly adjacent to u, we conclude that u cannot be the terminal b'_0 (i.e. $u = b'_i$ for $i > 1$). Therefore $b'_1 \in K_j$ for some $j < \mathtt{first}(b_{d-4})$; then by Proposition 8.5 and $b'_0 \sim b'_1$, it can be inferred that $b'_0 \notin ST(G)$. On the other hand, $b'_0 \not\sim c'_2$ implies $\mathtt{last}(b'_0) < \mathtt{last}(b_1)$; $\mathtt{last}(b'_{d'+1}) \not\sim c'_1$ implies $\mathtt{first}(b'_{d'}) > \mathtt{first}(b_d)$. Therefore, the base B' contains a b'_0-$b'_{d'}$ path and has to go through X (it cannot go through any vertex of N, which is adjacent all neighbors of u), which is impossible. ☐

Henceforth, therefore, we assume $c'_2 \notin N$.

Claim 3. $u = b'_0$.

Proof. According to Proposition 8.5, $\mathtt{last}(u) < \mathtt{first}(b_{d-2})$, and $N[u] \subseteq U_{0,\mathtt{first}(b_{d-2})}$. By Proposition 8.5 and Claims 1 and 2, $c'_2 \notin N[u]$, i.e. $c'_2 \not\sim u$, and thus u must be the terminal b'_0 ($u = b'_{d'+1}$ will contradict the selection of T). ☐

Claim 4. $b'_1 \in N$.

Proof. Recall $b'_0 = u$ and $N[u] \subseteq U_{0,\texttt{first}(b_{d-2})}$. Assume the contrary and $b'_1 \notin N$, then $c'_2 \not\sim b_d$, otherwise there is an AT with a shorter base than B. Observing $c'_2 \sim c_1, c_2$ as $c'_2 \in \texttt{seg}(T)$ and $c'_2 \not\sim s$ by Proposition 8.5, either $(s : c_1, c_2 : b_0, \dots b_\ell b'_0(u) b'_1 c'_2 b_d, b_{d+1})$ or $(s : c_1, c_2 : b_0, \dots b_\ell b'_0(u) b'_1 c'_2, b_{d+1})$ is such an AT, where $\ell \leqslant d - 5$ and b_ℓ is the last neighbor of b'_0 in B. From the facts $b'_1 \in U_{0,\texttt{first}(b_{d-2})} \setminus N$, $c'_2 \sim b'_1$ and $c'_2 \not\sim b_d$, in can be inferred that $\texttt{first}(b'_{d'}) < \texttt{last}(c'_2) < \texttt{first}(b_d)$, which, nonetheless, contradicts the selection of T. Therefore, $b'_1 \in N$. ☐

By Proposition 8.5 and $b'_1 \not\sim b'_3$, we conclude $\texttt{first}(b'_2) > \texttt{last}(b_{d-2})$. Now consider any vertex $x \in X \backslash Q$. If T' is a †-AT, then $N_{T'}(x) = \{b'_1\}$. If T' is a ‡-AT, then by Proposition 8.5 and Claim 1, $c'_1 \in N$; hence $N_{T'}(x) = \{b'_1, c'_1\}$. In either case, $T' \setminus \{u\} \cup \{x\}$ remains an AT. As this AT is contained in $G - Q$, we end with a contradiction. Therefore, Q' must be an interval deletion set of G. Now it remains to show the interval deletion set Q' we construct is minimum, from which the lemma follows.

Claim 5. $|Q'| \leqslant |Q|$.

Proof. By construction of Q' and the definition of X, it suffices to show that $Q \cap V_I$ makes a b_1-b_{d-4} separator in $G - N$; the claim ensues as $|Q'| = |Q \setminus V_I| + |X| \leqslant |Q \setminus V_I| + |Q \cap V_I| = |Q|$. Suppose the contrary and P is a chordless b_1-b_{d-4} path, then within the b_0-b_{d+1} path $P b_{d-3} b_{d-2} b_{d-1} b_d b_{d+1}$, we can construct a chordless b_0-b_{d+1} path P'. By Proposition 8.5, no vertex of P is adjacent to b_{d-1}, b_d, b_{d+1}; hence there are at least three vertices of P' are adjacent to c_1, c_2, but none is adjacent to s. As $s, c_1, c_2 \notin Q$ and P is disjoint from Q, we obtain an AT $(s : c_1, c_2 : l', B', r')$, where $B' \subseteq P'$, in $G - Q$, which is impossible. ☐

☐

As a matter of fact, if we are set to remove any vertex v, we actually remove all vertices in the nontrivial module that contains v. In particular, the set X specified in Lemma 8.6 is a nontrivial module.

Finding the AT required by Lemma 8.6 and branching on the deletion of one the 10 vertices completes the proof of Theorem 2.4.

Proof (of Theorem 2.4). Based on Lemma 8.6, it suffices to show how to find such an AT, and then the standard branching will deliver the claimed algorithm. For each seven vertices, we can check whether or not there is an AT T such that they are $\{s, c_1, c_2, b_0, b_1, b_d, b_{d+1}\}$ of T. Therefore, in $\mathcal{O}(n^8)$ we can check all combinations. On the other hand, since there are at most n nodes in the central path, there is a quadratic bound on the number of minimal segments. Now that we have such a minimal segment, the short base B can be easily found and thus we obtain the AT required by Lemma 8.6. ☐

9 Concluding Remarks

We have presented an $n^{\mathcal{O}(1)} \cdot 10^k$ algorithm for INTERVAL DELETION, and thus classified it to be FPT. The runtime of our algorithm should be contrasted with the currently best algorithm of runtime $n^{\mathcal{O}(1)} \cdot k^{2k}$ for INTERVAL COMPLETION [22], which, as common sense holds, should be easier than INTERVAL DELETION. Whereupon it is nature to ask the existence of an $n^{\mathcal{O}(1)} \cdot c^k$ time algorithm for INTERVAL COMPLETION; in particular, whether or not our approach can be applied to INTERVAL COMPLETION in some way? Another interesting comparison is with the CHORDAL DELETION, which still lacks a constructive FPT algorithm.

For INTERVAL DELETION, the constant $c = 10$ in the runtime $n^{\mathcal{O}(1)} \cdot c^k$ of our algorithm might be improvable. The constant 10 is determined by two branching steps that each branch into up to 10 instances. Let us briefly discuss the possibility of improving them. The first branching is on the small forbidden induced subgraphs where we destroy them in the simplest way. This might be improved using more careful manipulation and analysis. The other branching is conducted on asteroidal triples, where the graph is already chordal. As we have a linear organization of the central path and all asteroidal triples, dynamic programming might help here. Can it be improved to $c = 2$?

It is known that INTERVAL COMPLETION can be solved in polynomial time if the input graph is a circular-arc graph [16]. It would be interesting to inquire the complexity of INTERVAL DELETION and INTERVAL COMPLETION when the input graph is restricted to chordal graphs. Or at least, can it be solved in polynomial time if the input graph contains neither small asteroidal triples nor holes and every shallow terminal in it is simplicial, hence greatly improving Theorem 2.4. This, if positively answered, will imply that all the troubles are small forbidden subgraphs. We also leave it open for the INTERVAL EDGE DELETION problem, which instead asks for a set of k edges whose removal makes an interval graph. The obstacle here lies in the phase of breaking hole, where we no longer have a good bound for the number of edge hole covers for congenial holes.

References

[1] Isolde Adler, Martin Grohe, and Stephan Kreutzer. Computing excluded minors. In *SODA*, pages 641–650, 2008.

[2] Amotz Bar-Noy, Reuven Bar-Yehuda, Ari Freund, Joseph Naor, and Baruch Schieber. A unified approach to approximating resource allocation and scheduling. *Journal of the ACM*, 48(5):1069–1090, 2001.

[3] Seymour Benzer. On the topology of the genetic fine structure. *Proceedings of the National Academy of Sciences*, 45(11):1607–1620, 1959.

[4] Andreas Brandstädt, Van Bang Le, and Jeremy P. Spinrad. *Graph classes: a survey*. Society for Industrial and Applied Mathematics, Philadelphia, PA, USA, 1999.

[5] Peter Buneman. A characterization of rigid circuit graphs. *Discrete Mathematics*, 9(4):205–212, 1974.

[6] Leizhen Cai. Fixed-parameter tractability of graph modification problems for hereditary properties. *Information Processing Letters*, 58(4):171–176, 1996.

[7] Jianer Chen, Yang Liu, Songjian Lu, Barry O'Sullivan, and Igor Razgon. A fixed-parameter algorithm for the directed feedback vertex set problem. *Journal of the ACM*, 55(5), 2008.

[8] Gabriel A. Dirac. On rigid circuit graphs. *Abhandlungen aus dem Mathematischen Seminar der Universitt Hamburg*, 25(1):71–76, 1961.

[9] Rodney G. Downey and Michael R. Fellows. *Parameterized Complexity*. Springer, 1999.

[10] Fedor V. Fomin, Daniel Lokshtanov, Neeldhara Misra, and Saket Saurabh. Planar F-deletion: Approximation and optimal fpt algorithms, 2012. Accepted to FOCS 2012.

[11] Delbert R. Fulkerson and Oliver A. Gross. Incidence matrices and interval graphs. *Pacific Journal of Mathematics*, 15(3):835–855, 1965.

[12] Martin Charles Golumbic. *Algorithmic Graph Theory and Perfect Graphs*, volume 57 of *Annals of Discrete Mathematics*. North-Holland Publishing Co., Amsterdam, The Netherlands, 2004.

[13] Luciano Noberto Grippo and Martín Darío Safe. On circular-arc graphs having a model with no three arcs covering the circle. CLAIO-SBPO 2012, Rio de Janeiro - Brazil, September, 24-28 2012. http://www2.claiosbpo2012.iltc.br/pdf/102012.pdf.

[14] Haim Kaplan, Ron Shamir, and Robert Endre Tarjan. Tractability of parameterized completion problems on chordal, strongly chordal, and proper interval graphs. *SIAM Journal on Computing*, 28(5):1906–1922, 1999.

[15] Ken-ichi Kawarabayashi and Bruce A. Reed. An (almost) linear time algorithm for odd cyles transversal. In Yuval Rabani, editor, *SODA*, pages 365–378. SIAM, 2010.

[16] Ton Kloks, Dieter Kratsch, and C. K. Wong. Minimum fill-in on circle and circular-arc graphs. *J. Algorithms*, 28(2):272–289, 1998.

[17] C. G. Lekkerkerker and J. Ch. Boland. Representation of a finite graph by a set of intervals on the real line. *Fund. Math. Polska Akad. Nauk.*, 51:45–64, 1962.

[18] John M. Lewis and Mihalis Yannakakis. The node-deletion problem for hereditary properties is NP-complete. *Journal of Computer and System Sciences*, 20(2):219–230, 1980.

[19] Dániel Marx. Chordal deletion is fixed-parameter tractable. *Algorithmica*, 57(4):747–768, 2010.

[20] Dániel Marx and Ildikó Schlotter. Obtaining a planar graph by vertex deletion. *Algorithmica*, 62(3-4):807–822, 2012.

[21] Bruce A. Reed, Kaleigh Smith, and Adrian Vetta. Finding odd cycle transversals. *Operations Research Letters*, 32(4):299–301, 2004.

[22] Yngve Villanger, Pinar Heggernes, Christophe Paul, and Jan Arne Telle. Interval completion is fixed parameter tractable. *SIAM Journal on Computing*, 38(5):2007–2020, 2009.

Appendix: Examples Used in Proofs



(a) $x \sim B$ (b) $x \not\sim B$ but $x \sim \{l, r\}$



(c) $x \sim s$ but $x \not\sim B$ (d) $x \sim s$ but $x \not\sim B$ (e) $x \sim c_1$ but $x \not\sim c_2, l$

Figure 6: A neighbor (x) of the shallow terminal that is not adjacent to the center (c or c_2)
Lemma 5.1.



(a) †-AT, $x \sim l, r$ (b) †-AT, $x \not\sim l$, and $x \sim r$ (c) †-AT, $x \not\sim l, r$



(d) ‡-AT, $x \sim l, r$ (e) ‡-AT, $x \not\sim l$, and $x \sim r$ (f) ‡-AT, $x \not\sim l, r$

Figure 7: A common neighbor x of B that is not adjacent to a neighbor v of s
Lemma 5.1.



(a) $x \sim l$.



(b) $x \sim b_1$ but $x \not\sim l$.



(c) $x \not\sim l, b_1$.

Figure 8: (x) is adjacent to both the shallow terminal (s) and the base B
Lemma 5.1.



(a) $T_1 = (s : c : l, B, r)$
$T_2 = (r : b_d : s, B', r')$.

(b) $T_1 = (s : c_1, c_2 : l, B, r)$
$T_2 = (r : c_2, b_d : s, B', r')$.

NOTES

- s can be replaced by any b_i $(i \leqslant d-3)$ in T_2;
- similarly, r can be replaced by any b'_i $(i \geqslant 4)$ in T_1;
- In (b), $(r : c_2 : l, B', r')$ also makes an †-AT.

Figure 9: Interaction between ATs
T_1 edges in black, T_2 edges in blue, and shared edges in red.